

540,759  
Rec'd PCT/PTO 24 JUN 2005

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



10/540759



(43) International Publication Date  
15 July 2004 (15.07.2004)

PCT

(10) International Publication Number  
WO 2004/058814 A1

(51) International Patent Classification<sup>7</sup>: C07K 14/415,  
C07H 21/04, C12N 15/63, 15/82

Albert Road, Mt Albert (NZ). MATICH, Adam [AU/NZ];  
HortResearch, Tennent Drive, Palmerston North (NZ).

(21) International Application Number:  
PCT/NZ2003/000294

(74) Agents: ADAMS, Matthew, D., et al.; A J Park, 6th Floor  
Huddart Parker Building, Post Office Square, PO Box 949,  
6015 Wellington (NZ).

(22) International Filing Date:  
24 December 2003 (24.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
523384 24 December 2002 (24.12.2002) NZ

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,  
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,  
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,  
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,  
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,  
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): THE  
HORTICULTURE AND FOOD RESEARCH INSTI-  
TUTE OF NEW ZEALAND LIMITED [NZ/NZ]; Mt  
Albert Research Centre, 120 Mt Albert Road, Mt Albert  
(NZ).

(84) Designated States (*regional*): ARIPO patent (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,  
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): GREEN, Sol  
[NZ/NZ]; 3/3 Park Ave, Takapuna (NZ). FRIEL, Ellen  
[GB/NZ]; 8/23B Western Springs Road, Western Springs,  
Auckland (NZ). BEUNING, Lesley [NZ/NZ]; 11 Meryl  
Avenue, Huapai (NZ). MACRAE, Elspeth [NZ/NZ];  
The Horticulture and Food Research Institute of New  
Zealand Limited, Mt Albert Research Centre, 120 Mt

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: ENZYMES AND POLYNUCLEOTIDES ENCODING THE SAME

(57) Abstract: The invention relates to a multifunctional germacrene-D synthase and a polynucleotide encoding it. Included also are genetic constructs, transgenic organisms, plant selection methods and biofermentation methods. The enzyme may be used to prepare germacrene-D and/or other sesquiterpenes such as germacrene-D, delta-cadinene, gamma-cadinene, gamma-murolene, gamma-elemene, delta-elemene, elemol or germacrene-B.

WO 2004/058814 A1

## ENZYMES AND POLYNUCLEOTIDES ENCODING THE SAME

### Technical Field

5 The present invention relates to the enzyme multifunctional germacrene-D synthase and to polynucleotide sequences encoding the enzyme. The invention also relates to nucleic acid constructs, vectors and host cells incorporating the polynucleotide sequences. It further relates to the production of a multifunctional germacrene-D synthase and more specifically to the production of germacrene-D, *gamma*-muurolene and *delta*- and *gamma*-cadinene and minor quantities of other sesquiterpenes and their use in products  
10 such as an insect attractant/deterrent, anti-bacterial and anti-fungal agents, flavours and fragrances and other products. Germacrene-D, *gamma* muurolene and *delta*- and *gamma*-cadinene may also be used to produce further products with characteristic aromas useful as flavours and fragrances, or products that are useful as insect  
15 attractants/deterrents or antimicrobial agents.

### Background Art

Germacrene-D (Figure 1) is a cyclic sesquiterpene hydrocarbon (C<sub>15</sub>H<sub>24</sub>; 7-iso-propyl-  
20 10-methyl-4-methylene-cyclodeca-5,10-diene) that is either constitutively present or induced in a wide range of plant species. *Delta*-Cadinene, *gamma*-muurolene and *gamma*-cadinene (Figure 1) are cyclic sesquiterpene hydrocarbons that are either constitutively present or induced in a wide range of plant species.

25 The biosynthetic pathway for the sesquiterpenes branches off from the general terpenoid pathway, beginning with the allylic diphosphate ester farnesyl diphosphate (FDP, also called FPP) (Bohlmann, *et al.*, Proc. Natl. Acad. Sci. U. S. A. 95, 4126-4133 (1998), Cane and Bowser, Bioorg. Med. Chem. Lett. 9, 1127-1132 (1999), Davis and Croteau, Top. Curr. Chem 209, 53-95 (2000)). Germacrene-D and *delta*-Cadinene can both be  
30 synthesised from FDP in a reaction that proceeds through a carbocation intermediate (Figure 2) and are catalysed by the sesquiterpene synthases germacrene-D synthase (Guterman *et al* Plant Cell 14, 2325-2338 (2002)) and *delta*-cadinene synthase

(Benedict *et al* Plant Phys 125, 1754-1765 (2001)) respectively. No gene has been isolated that has been demonstrated to produce significant quantities of gamma-muurolene or gamma cadinene. The pathway for sesquiterpene biosynthesis, the acetate/mevalonate pathway, is localised to the cytoplasm; in contrast to the pathways for monoterpene and diterpene biosynthesis, which occur in the chloroplast (Lange, *et al.*, Proc. Natl. Acad. Sci. U. S. A 97, 13172-13177 (2000)). However recent labelling studies have shown that germacrene D in *Solidago canadensis* is formed predominantly via the methylerythritol phosphate pathway (Steliopoulos *et al.*, Phytochemistry 60: 13-20 (2002)).

All known plant terpene synthases, however, whether monoterpene, sesquiterpene or diterpene, appear to be closely related. Similarities include the positioning of intron sequences (Trapp and Croteau, Genetics 158, 811-832 (2001)) and the presence of conserved sequences, such as an aspartate-rich DDX(X,D,E) motif (Lesburg, *et al.*, 8, 695-703 (1998)). This motif is involved in the binding of metal ions, usually  $Mg^{2+}$ , that are necessary for catalysis.

Germacrene-D is considered to be a key intermediate in the biosynthesis of many sesquiterpenes (Yoshihara *et al.*, Tetrahed Lett 27: 2263-2364 (1969); Bülow and König, Phytochem 55, 141-168 (2000)). Furthermore, it has been shown to increase attraction of and oviposition by the tobacco budworm moth *Heliothis virescens* (Mozuraitis *et al* Chem Senses 27, 505-509 (2002)) and is also a sex stimulant for the male American cockroach (*Periplaneta americana* L.) (Nishino *et al.*, J Insect Physiol 23, 415-419 (1977)). Germacrene-D, as a component of certain essential oils, has also been shown to possess antibacterial properties (Juteau *et al* Fitoterapia 73, 532-535 (2002)). Germacrene D is also a major volatile in leaves of *Solidago* species (Prosser *et al.* Phytochemistry 60, 691-702 (2002)) and in lemon basil (Iijima *et al.* Plant Physiology 134, 1-10 (2004)). In general the (R)-(+)-enantiomer is found in lower plants, whereas the (S)-(-) enantiomer is found in higher plants, although there are plants where both enantiomers exist (Schmidt *et al.*, Angewandte Chemie 110: 1479-1481 (1998); Schmidt *et al.*, Chirality 11:353-362 (1999)). To date a gene for germacrene-D synthase has been isolated only from rose (Guterman *et al* Plant Cell 14,

2325-2338 (2002)), which appeared to produce germacrene-D only. Its enantiomeric form is unknown.

5 Cadinene is the first intermediate in the conversion by *delta*-cadinene synthase of FDP to sesquiterpene phytoalexins in cotton (*Gossypium barbadense*) (Benedict *et al* Plant Phys 125, 1754-1765 (2001); Davis and Essenberg, Phytochem 39, 553-567 (1995); Davis *et al.*, Phytochem 41, 1047-1055 (1996)), and is the precursor of deoxyhemigossypol and hemigossypol defense sesquiterpenes. *Alpha*-cadinol has been shown to be a potent miticide against house dust mites (Chang *et al.*, J. Med Entom. 10 38:455-458 (2001)). To date four cDNAs for *delta*-cadinene synthase have been isolated from *Gossypium arboreum* and characterisation of at least one of these has been reported (Chen *et al*, Arch. Biochem Biophys 324, 255-266 (1995); Meng *et al*, J. Nat. Prod. 62, 248-252 (1999)). Although *gamma*-cadinene and *gamma*-muurolene are present widely in plants, no genes encoding for proteins that allow production of any of 15 these compounds in any quantity have been discovered.

The applicants have identified a polynucleotide encoding a multifunctional germacrene-D synthase which facilitates the production of germacrene-D, *gamma*-muurolene and 20 *delta*- and *gamma*-cadinene in biofermentation processes. By manipulation of the process conditions, the applicant obtained germacrene-D or germacrene-D with a mix of *delta*-cadinene, *gamma*-cadinene and *gamma*-muurolene in different ratios. Minor amounts of *delta*-elemene, germacrene B, elemol and *delta*-cadinene may be coproduced with germacrene-D. Minor amounts of *alpha*-cubebene, *alpha*-ylangene, 25 *alpha*-copaene, *beta*-cubebene, isodene, epibicyclosesquiphellandrene and *alpha*-cadinol/*beta*-eudesmol may be coproduced under acidic conditions with major quantities of *gamma*-muurolene, *delta*-cadinene, *gamma*-cadinene and germacrene-D. The polynucleotide can also be used to co-ordinately manipulate production of germacrene-D, *delta*-elemene, germacrene-B, elemol and *delta*-cadinene in transgenic 30 plants to alter fragrance/flavour characteristics and/or plant pathogen and/or insect interactions. The gene can also be used as a marker in marker assisted breeding to



discover plant material with altered germacrene-D, and *delta*-cadinene, *delta*-elemene, elemol or germacrene-B composition.

### Summary of the Invention

5

In a first aspect the invention provides an isolated polynucleotide encoding a multifunctional germacrene-D synthase.

10

In a further aspect the invention provides an isolated polynucleotide comprising the sequence shown in Figure 3 (SEQ ID NO:1) or a fragment or variant thereof encoding a polypeptide with multifunctional germacrene-D synthase activity.

15

The polypeptides of the invention are multifunctional. They are capable of facilitating the conversion of FDP to a mixture of germacrene-D and *gamma*-muurolene, *delta*-and *gamma*-cadinene and other minor sesquiterpenes. Depending on the manner of trapping of the volatile products, the recombinant enzyme, can be used to obtain: germacrene D (plus minor amounts of other sesquiterpenes), or germacrene D with a mix of *delta*-and *gamma*-cadinene and *gamma*-muurolene (plus minor amounts of other sesquiterpenes)

20

In a further aspect, the invention provides an isolated polynucleotide encoding the polypeptide shown in Figure 4 (SEQ ID NO:2) or encoding a variant or a fragment of that sequence which has a multifunctional germacrene-D synthase activity.

25

In a further aspect the invention provides an isolated multifunctional germacrene-D synthase polypeptide.

30

In yet a further aspect, the invention provides an isolated multifunctional germacrene-D synthase having the sequence shown in Figure 4 (SEQ ID NO:2) or a fragment or variant thereof with similar multifunctional germacrene-D synthase activity.

The polypeptides of the invention are useful for *in vitro* preparation of germacrene-D and/or *delta*-cadinene, and/or *gamma*-muurolene, and/or *gamma*-cadinene. They may

also be used for *in vitro* preparation of *delta*-elemene, germacrene B, *gamma*-elemene or elemol, or *alpha*-cubebene, *alpha*-ylangene, *alpha*-copaene, *beta*-cubebene, isodene, epibicyclosesquiphellandrene and *alpha*-cadinol/*beta*-eudesmol.

- 5 In a further aspect the invention provides a vector comprising a polynucleotide of the invention.

In a further aspect the invention provides a genetic construct comprising an open reading frame polynucleotide encoding a polypeptide of the invention.

10

In yet a further aspect the invention provides a genetic construct comprising in the 5'-3' direction

- (a) a promoter sequence; and
- (b) an open reading frame polynucleotide encoding a polypeptide of the invention

15

Preferably the genetic construct also comprises a termination sequence.

In another aspect the invention provides a genetic construct comprising in the 5'-3' direction

20

- (a) a promoter sequence; and
- (b) a polynucleotide which hybridizes to a polynucleotide encoding a polypeptide of the invention

Preferably the genetic construct also comprises a termination sequence.

- 25 In a further aspect the invention provides a host cell comprising a genetic construct of the invention.

In still a further aspect, the invention provides a transgenic plant cell which includes a genetic construct of the invention.

30

In a yet further aspect, the invention provides a plant cell which has been modified to alter expression of a multifunctional germacrene-D synthase.

In addition the invention provides a transgenic plant comprising such cells.

In another aspect the invention provides a method of preparing germacrene-D, *delta*-cadinene, *gamma*-cadinene, *gamma*-muurolene, *gamma*-elemene, *delta*-elemene, elemol or germacrene B comprising the steps of

- (a) culturing a cell which has been genetically modified with a polynucleotide of the invention to provide increased multifunctional germacrene-D synthase activity;
- (b) providing the cell with farnesyl diphosphate or geranyl diphosphate if necessary; and
- (c) separating the germacrene-D and/or *delta*-cadinene and/or and/or *delta* elemene and/or elemol and/or germacrene B, and/or *gamma*-cadinene, and/or *gamma*-muurolene, and/or *gamma*-elemene produced.

This method of the invention allows use of biofermentation as a convenient method for preparing the product.

Depending on the manner of trapping of the volatile products from the recombinant enzyme, it is possible to obtain germacrene D, elemol, *delta*-elemene, germacrene B and *delta*-cadinene, or germacrene D with a mix of *delta* cadinene, *gamma* muurolene and *gamma*-cadinene (plus minor amounts of other sesquiterpenes)

This gives us the opportunity to produce all these major compounds with the one protein. By varying the conditions in non plant production systems each compound may be obtained in a controlled manner.

In further aspect the invention provides a method for modulating the Germacrene-D and/or *delta*-cadinene and/or germacrene B and or elemol and/or *delta*-elemene, and/or *gamma*-cadinene, and/or *gamma*-muurolene, and/or *gamma*-elemene production of a plant, the method comprising: increasing or decreasing expression of multifunctional germacrene-D synthase wherein said increasing or decreasing is achieved by genetic

modification to alter the expression of a gene encoding a multifunctional germacrene-D synthase.

In a further aspect the invention provides a method for modulating germacrene-D and/or *delta*-cadinene and/or germacrene B and or elemol and/or *delta*-elemene, and/or *gamma*-cadinene, and/or *gamma*-muurolene, and/or *gamma*-elemene production, in a plant, the method comprising of :

- (a) introducing into the plant, a genetic construct of the invention; and
- (b) transcriptionally expressing the polynucleotide in the plant.

In a further aspect the invention provides a method for modulating a multifunctional germacrene-D synthase production in a plant, the method comprising of

- (a) introducing into the plant, a DNA genetic construct of the invention; and
- (b) expressing a polypeptide of the invention in the plant.

In a further aspect the invention provides a method of selecting a plant with altered germacrene-D (and/or or other sesquiterpenes as above) content comprising the steps of:

- (a) contacting polynucleotides from at least one plant with at least one polynucleotide comprising at least 15 contiguous nucleotides of the polynucleotide of the invention to assess the expression of the multifunctional germacrene-D synthase; and
- (b) selecting a plant showing altered expression.

Preferably the plant is an angiosperm, more preferably selected from Actinidiaceae and Ericaceae, most preferably from Actinidiaceae.

In yet a further aspect the invention provides a method for preparing germacrene-D and/or other sesquiterpenes comprising:

- (a) obtaining a polypeptide of the invention; and
- (b) incubating farnesyl diphosphate in the presence of the polypeptide, and
- (c) separating the germacrene D and and/or other compounds produced.

.In a further aspect the invention provides an isolated polynucleotide comprising a sequence selected from the sequences of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4 and SEQ ID NO: 7 and SEQ ID NO: 9, or a fragment or variant thereof encoding a polypeptide with multifunctional germacrene-D synthase activity.

5

In a further aspect the invention provides an isolated polypeptide with multifunctional germacrene-D synthase activity comprising a sequence selected from SEQ ID NO: 2, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10, or a fragment or variant thereof.

10

### **Brief Description of Drawings**

The present invention will be better understood with reference to the accompanying drawings in which:

15

Figure 1 shows the structure of Germacrene D and other sesquiterpenes produced by EST 75565.

20

Figure 2 shows (A) a pathway for the synthesis of germacrene D from FDP and (B) examples of acid catalysed rearrangement products of germacrene D.

25

Figure 3 shows the polynucleotide sequence that encodes a multifunctional germacrene-D synthase. The sequence was obtained from a cDNA library that was constructed using mRNA extracted from *Actinidia deliciosa* petals.

Figure 4 shows the predicted amino acid sequence of a multifunctional germacrene-D synthase from *A. deliciosa* petals.

30

Figure 5 shows the polynucleotide sequences of ESTs 72838, 80968, 304951 and 82293.

Figure 6 shows the amino acid sequence encoded by ESTs 72838, 80968, 304951 and 82293.

5 Figure 7 shows the presence (a: total ion chromatogram, b: ion 161 ) of Germacrene D from the headspace volatiles in *Actinidia deliciosa* flowers.

Figure 8 shows a GC-MS trace of headspace above partially purified cell free extracts (optimum activity conditions) showing internal standard (C16) and sesquiterpene trapped on Carbowax®/DVB SPME fibres after 21 hours

10

Figure 9 shows a GC-MS trace of headspace above partially purified cell free extracts (optimum activity conditions).showing internal standard (C16) and sesquiterpene products trapped on Chromosorb 105 cartridge after 18 hours.

15 Figure 10 shows a GC-MS trace of headspace above bacterial cultures harbouring (a) pET control vector or (b) pET 30/75565. FDP was added to both cultures. Trapping used PDMS/DVB fibres.

20 Figure 11 shows the effect of trapping matrix on sesquiterpene composition, in particular sesquiterpene products from pET30/75565 purified protein extract + FDP trapped on PDMS/DVB SPME fibres after 24 hours.

25 Figure 12 shows the results of a time course experiment showing how the ratio of Germacrene D changes relative to the other Germacrene D derivative compounds over time A) at t=0 after addition of FDP, B) at t=1 hr after addition of FDP and C) at t=2 hrs after addition of FDP.

30 Figure 13 shows the change in headspace volatile profile trapped on PDMS/DVB fibres over a time course of 24 hours.

Figure 14 shows graphs of the variation of the multifunctional germacrene-D synthase activity with (A) substrate concentration, (B)  $Mg^{2+}$  concentration (C)  $Mn^{2+}$  concentration and (D) pH.

- 5 Figure 15 shows graph of germacrene-D synthase activity against temperature.

Figure 16 shows variation of germacrene-D synthase activity with varying metal ion contents.

- 10 Figure 17 shows GC-MS trace of headspace profiles at 3 different pHs trapped on Carbowax®/DVB SPME fibres after 21 hours.

Figure 18 shows Northern and RT-PCR analysis of total RNA extracted from different tissues of *Actinidia deliciosa* and from *Actinidia chinensis* fruit. (A) Northern analysis

- 15 using as a probe an approximately 350 base pair [ $\alpha$ - $^{32}P$ ]dCTP-labelled PCR fragment that was amplified from EST 75565 with the primers 75565NF1 (5' TAGGCGTGTCCTTACCATTTT 3') (SEQ ID NO: 15) and 75565NR1 (5' GCATGAATGATTTGTTTCCTT 3') (SEQ ID NO: 16). (B) RT-PCR analysis using two sets of PCR primers that amplify an approximately 210 base pair fragment about  
20 1000 base pairs in from the start ATG (Primer set 1) and an approximately 350 base pair fragment about 450 base pairs in from the start ATG (Primer set 2). -ve = PCR amplification using Primer set 1.

- Figure 19 shows RT-PCR amplification of total RNA extracted from seedlings of transgenic *Arabidopsis thaliana* lines using primers (Primer set 2) designed towards the  
25 5' end of the germacrene D synthase cDNA sequence. The size of the amplification product expected is approximately 350 bp. The lane labelled MW is the molecular weight marker (Invitrogen). Lanes labelled Line 1 to 4 contain the resulting products from PCR amplification of the total RNA and Lanes labelled 2 contain the resulting products from RT-PCR amplification of the total RNA. bp=base pairs.

Figure 20 Profiles of ions (A) 161 and (B) 204 (indicative of sesquiterpenes) from the GCMS trace of headspace collected above *Arabidopsis* plants transformed with multifunctional germacrene D synthase gene growing in a 3L growth vessel for 6 days.

## 5 Detailed Description

In one embodiment of the invention, cells genetically modified to exhibit a multifunctional germacrene-D synthase activity are used for the production of germacrene-D and/or *delta*-cadinene, and/or elemol and/or *delta*-elemene and/or germacrene B. While the cells may potentially be of any cell type that can be grown in culture, it is currently preferred to use bacteria or yeast cells for producing germacrene-D and/or *delta*-cadinene and other compounds (and oxidation products or derivatives). Preferred cells for use in the biofermentation processes of this embodiment are GRAS microbes; eg *E. coli*, *Lactobacillus* spp and other non-pathogenic bacteria or yeasts such as brewers yeast or fungi such as *Aspergillus niger*. Methods of capture of the produced compounds will be important in determining the composition eg. materials such as Chromasorb results predominantly in germacrene D, whereas silica-based matrices will result in predominantly *gamma*-muurolene and *delta*-cadinene.

Germacrene-D and its co- or by- products may be used as flavour and fragrance additives, an antimicrobial agent, as a pheromone for attracting or repelling insects. Delta-cadinene and its co- or by- products may be used as flavour and fragrance additives (eg baking goods, detergents, cosmetics, chewing gum), and as a pheromone repellent, particularly for cotton. Both compounds may be used as precursors for synthesis of other compounds of value.

In another aspect of the invention, the polynucleotides of the invention are used to prepare transgenic plants that over-express the multifunctional germacrene-D synthase in at least some parts of the plant. In this way the invention is used to impart fragrance to flowers, repel or attract insects (either as indicator plants, host plants, or alternative



hosts) or impart an altered flavour to fruit or prevent disease in fruit, or to extract pharmaceutical products or animal or insect efficacious extracts.

In one particular aspect the polynucleotides of the invention are used in plants of the order Actinidiaceae and Ericaceae, particularly in the genus *Actinidia* and the genus *Vaccinium* to provide increased fragrance in flowers (or fruit). This aspect has particular use in kiwifruit and blueberries.

In another aspect polynucleotides of the invention are used to decrease multifunctional germacrene-D synthase activity in kiwi fruit. This may be achieved in several ways, for example by genetically modifying the kiwi fruit so that the multifunctional germacrene-D synthase polynucleotide is transcribed in an antisense orientation which results in decreased multifunctional germacrene-D synthase translation. Such fruit may then be more resistant to certain diseases and pests.

Altering the levels of germacrene-D and/or *delta*-cadinene and/or other components in flowers may alter attraction/repulsion of pollinators or may also affect invasion by pathogens.

In another aspect the invention provides a method useful in kiwifruit or blueberry breeding. Segments of the polynucleotide sequences of the invention may be used as probes to investigate the genetic makeup of candidate kiwifruit or blueberry varieties with respect to multifunctional germacrene-D synthase activity. The presence of high levels of polynucleotides encoding multifunctional germacrene-D synthase activity in the flowers or fruit of kiwifruit or blueberry may be used to identify kiwifruit or blueberries with altered fragrance or flavours.

The amino acid sequence of one polypeptide, a multifunctional germacrene-D synthase from kiwifruit, and that of the polynucleotide sequence encoding it are given in Figures 4 and 3 respectively (SEQ ID NO:2 and SEQ ID NO:1). It will however be appreciated

that the invention is not restricted only to the polynucleotide/polypeptide having the specific nucleotide/amino acid sequence given in Figures 3 and 4. Instead, the invention also extends to variants of the polynucleotide/polypeptide of Figures 3 and 4 which possess or encode a multifunctional germacrene-D synthase activity.

5

The term "polynucleotide(s)" as used herein means a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases and includes DNA and corresponding RNA molecules, including hnRNA and mRNA molecules, both sense and anti-sense strands, and comprehends cDNA, genomic DNA and recombinant DNA, as well as wholly or partially synthesized polynucleotides. An hnRNA molecule contains introns and corresponds to a DNA molecule in a generally one-to-one manner. An mRNA molecule corresponds to an hnRNA and DNA molecule from which the introns have been excised. A polynucleotide may consist of an entire gene, or any portion thereof. Operable anti-sense polynucleotides may comprise a fragment of the corresponding polynucleotide, and the definition of "polynucleotide" therefore includes all such operable anti-sense fragments.

10

15

The term 'polypeptide(s)' as used herein includes peptides, polypeptides and proteins.

The phrase "variants with multifunctional germacrene-D synthase activity" is used in recognition that it is possible to vary the amino acid/nucleotide sequence of a polypeptide/polynucleotide while retaining substantially equivalent functionality. The equivalent can be, for example, a fragment of the polypeptide, a fusion of the polypeptide with another polypeptide or carrier, or a fusion of a fragment with additional amino acids.

20

25

An "isolated" polypeptide is a polypeptide that has been identified and separated or recovered to be largely free of components of its natural environment, (that is so that the polypeptide comprises at least 50% of the polypeptides from its natural environment, preferably at least 80%, more preferably at least 90%). The term "isolated" polypeptide includes polypeptides in situ within recombinant cells. However generally isolated polypeptides will be prepared by at least one purification step.

30

An "isolated" polynucleotide is a nucleotide molecule that is identified and separated from at least one contaminant polynucleotide with which it is ordinarily associated.

- 5 Variant polynucleotide sequences also include equivalent sequences, which vary in size, composition, position and number of introns, as well as size and composition of untranslated terminal regions. Variant polynucleotides also include those encoding functionally equivalent polypeptides.
- 10 It will be understood that a variety of substitutions of amino acids is possible while preserving the structure responsible for activity of the polypeptides. Conservative substitutions are described in the patent literature, as for example, in United States Patent No 5,264,558 or 5,487,983. It is thus expected, for example, that interchange among non-polar aliphatic neutral amino acids, glycine, alanine, proline, valine and
- 15 isoleucine, would be possible. Likewise, substitutions among the polar aliphatic neutral amino acids, serine, threonine, methionine, asparagine and glutamine could be made. Substitutions among the charged acidic amino acids, aspartic acid and glutamic acid, could probably be made, as could substitutions among the charged basic amino acids, lysine and arginine. Substitutions among the aromatic amino acids, including
- 20 phenylalanine, histidine, tryptophan and tyrosine are also possible. Such substitutions and interchanges are well known to those skilled in the art.

Equally, nucleotide sequences encoding a particular product can vary significantly simply due to the degeneracy of the nucleic acid code.

25

- A polynucleotide or polypeptide sequence may be aligned, and the percentage of identical nucleotides in a specified region may be determined against another sequence, using computer algorithms that are publicly available. An exemplary algorithm for aligning and identifying the similarity of polynucleotide sequences is the BLASTN
- 30 algorithm. The similarity of polypeptide sequences may be examined using the BLASTP algorithm. Both the BLASTN and BLASTP software are available on the NCBI anonymous FTP server (<ftp://ncbi.nlm.nih.gov>) under `/blast/executables/`. The

BLASTN algorithm version 2.0.4 [Feb-24-1998], set to the default parameters described in the documentation of variants according to the present invention. The use of the BLAST family of algorithms, including BLASTN and BLASTP, is described at NCBI's website at URL <http://www.ncbi.nlm.nih.gov/BLAST/newblast.html> and in the publication of Altschul *et al.*, Nucleic Acids Res. 25, 3389-34023 (1997).

The following running parameters are preferred for determination of alignments and similarities using BLASTN that contribute to E values (as discussed below) and percentage identity: Unix running command: `blastall -p blastn -d embldb -e 10 -G 1 -E`

10 `1 -r 2 -v 50 -b 50 -I queryseq -o results`; and parameter default values:

`-p` Program Name [String]

`-d` Database [String]

`-e` Expectation value (E) [Real]

`-G` Cost to open a gap (zero invokes default behaviour) [Integer]

15 `-E` Cost to extend a cap (zero invokes default behaviour) [Integer]

`-r` Reward for a nucleotide match (blastn only) [Integer]

`-v` Number of one-line descriptions (V) [Integer]

`-b` Number of alignments to show (B) [Integer]

`-i` Query File [File In]

20 `-o` BLAST report Output File [File Out] Optional

For BLASTP the following running parameters are preferred: `blastall -p blastp -d swissprot -e 10 -G 1 -E 1 -v 50 -b 50 -I queryseq -o results`

`-p` Program Name [String]

`-d` Database [String]

25 `-e` Expectation value (E) [Real]

`-G` Cost to open a gap (zero invokes default behaviour) [Integer]

`-E` Cost to extend a cap (zero invokes default behaviour) [Integer]

`-v` Number of one-line descriptions (v) [Integer]

`-b` Number of alignments to show (b) [Integer]

30 `-i` Query File [File In]

`-o` BLAST report Output File [File Out] Optional

The "hits" to one or more database sequences by a queried sequence produced by BLASTN, BLASTP, or a similar algorithm, align and identify similar portions of sequences. The hits are arranged in order of the degree of similarity and the length of sequence overlap. Hits to a database sequence generally represent an overlap over only  
5 a fraction of the sequence length of the queried sequence.

The BLASTN algorithm also produces "Expect" or E values for alignments. The E value indicates the number of hits one can "expect" to see over a certain number of contiguous sequences by chance when searching a database of a certain size. The  
10 Expect value is used as a significance threshold for determining whether the hit to a database, such as the preferred EMBL database, indicates true similarity. For example, an E value of 0.1 assigned to a hit is interpreted as meaning that in a database of the size of the EMBL database, one might expect to see 0.1 matches over the aligned portion of the sequence with a similar score simply by chance. By this criterion, the aligned and  
15 matched portions of the sequences then have a 90% probability of being the same. For sequences having an E value of 0.01 or less over aligned and matched portions, the probability of finding a match by chance in the EMBL database is 1% or less using the BLASTN algorithm.

20 According to one embodiment, "variant" polynucleotides, with reference to each of the polynucleotides of the present invention, preferably comprise sequences having the same number or fewer nucleic acids than each of the polynucleotides of the present invention and producing an E value of 0.01 or less when compared to the polynucleotide of the present invention. That is, a variant polynucleotide is any  
25 sequence that has at least a 99% probability of being the same as the polynucleotide of the present invention, measured as having an E value of 0.01 or less using the BLASTN algorithm set at the parameters discussed above.

30 Variant polynucleotide sequences will generally hybridize to the recited polynucleotide sequence under stringent conditions. As used herein, "stringent conditions" refers to prewashing in a solution of 6X SSC, 0.2% SDS; hybridizing at 65°C, 6X SSC, 0.2% SDS overnight; followed by two washes of 30 minutes each in 1X SSC, 0.1% SDS at

65°C and two washes of 30 minutes each in 0.2X SSC, 0.1% SDS at 65°C. The variant polynucleotide sequences of the invention are at least 50 nucleotides in length.

5 Variant polynucleotides also include sequences which have a sequence identity of at least 60%, generally 70%, preferably 80%, more preferably 90%, even more preferably 95%, very preferably 98% and most preferably 99% or more to the nucleotide sequence given in the sequence listing herein.

10 In general, polypeptide sequences that code for the multifunctional germacrene-D synthases of the invention will be at least 60%, preferably 70%, and even 80%, 85%, 90%, 95%, 98%, most preferably 99% homologous or more with the disclosed amino acid sequence. That is, the sequence similarity may range from 60% to 99% or more. In addition the invention includes polynucleotide sequences encoding these amino acid sequences.

15 Also encompassed by the invention are fragments of the polynucleotide and polypeptide sequences of the invention. Polynucleotide fragments may encode protein fragments which retain the biological activity of the native protein. Alternatively, fragments used as hybridisation probes generally do not encode biologically active sequences.

20 Fragments of a polynucleotide may range from at least 15, 20, 30, 50, 100, 200, 400 or 1000 contiguous nucleotides up to the full length of the native polynucleotide sequences disclosed herein.

25 Fragments of the polypeptides of the invention will comprise at least 5, 10, 15, 30, 50, 75, 100, 150, 200, 400 or 500 contiguous amino acids, or up to the total number of amino acids in the full length polypeptides of the invention.

30 Variant is also intended to allow for rearrangement, shifting or swapping of one or more nucleotides or domains/motifs (from coding, non-coding or intron regions) from genes (including terpene synthases) from the same or other species, where such variants still provide a functionally equivalent protein or polypeptide of the invention or fragment thereof.

It is, of course, expressly contemplated that homologs to the specifically described multifunctional germacrene-D synthase having the sequence of Figure 4 (SEQ ID NO:2) exist in other plants. Such homologs are also "variants" as the phrase is used  
5 herein.

A polynucleotide sequence of the invention may further comprise one or more additional sequences encoding one or more additional polypeptides, or fragments thereof, so as to encode a fusion protein. Systems for such recombinant expression  
10 include, but are not limited to, mammalian, yeast, bacteria and insect systems.

DNA sequences from plants other than *A. deliciosa* which are homologs of the multifunctional germacrene-D synthase of Figure 3 (SEQ ID NO:1) may be identified (by computer-aided database searching) and isolated following high throughput  
15 sequencing of cDNA libraries prepared from such plants. Alternatively, oligonucleotide probes based on the sequence of Figure 4 (SEQ ID NO:2) can be synthesized and used to identify positive clones in either cDNA or genomic DNA libraries from other plants by means of hybridization or PCR techniques. Probes should be at least about 10, preferably at least about 15 and most preferably at least about 20 nucleotides in length.

20 Hybridization and PCR techniques suitable for use with such oligonucleotide probes are well known in the art. Positive clones may be analyzed by restriction enzyme digestion, DNA sequencing or the like.

The polynucleotides of the present invention may be generated by synthetic means  
25 using techniques well known in the art. Equipment for automated synthesis of oligonucleotides is commercially available from suppliers such as Perkin Elmer/Applied Biosystems Division (Foster City, CA) and may be operated according to the manufacturer's instructions.

30 As a result of the identification of the polypeptides and polynucleotides of the invention multifunctional germacrene-D synthase activity may be modulated in plants.

Modulation may involve a reduction in the expression and/or activity (i.e. silencing) of the polypeptide.

Any conventional technique for effecting this can be employed. Intervention can occur post-transcriptionally or pre-transcriptionally. Further, intervention can be focused upon the gene itself or on regulatory elements associated with the gene and which have an effect on expression of the encoded polypeptide. "Regulatory elements" is used here in the widest possible sense and includes other genes which interact with the gene of interest.

Pre-transcriptional intervention can involve mutation of the gene itself or of its regulatory elements. Such mutations can be point mutations, frameshift mutations, insertion mutations or deletion mutations. These latter mutations include so called "knock-out" mutations in which expression of the gene is entirely ablated.

Examples of post-transcriptional interventions include co-suppression or anti-sense strategies, a dominant negative approach, or techniques which involve ribozymes to digest, or otherwise be lethal to, RNA post-transcription of the target gene.

Co-suppression can be effected in a manner similar to that discussed, for example, by Napoli *et al.* *Plant Cell* 2, 279-290 (1990) and de Carvalho Niebel *et al.* *Plant Cell* 7, 347-358 (1995). In some cases, it can involve over-expression of the gene of interest through use of a constitutive promoter. It can also involve transformation of a plant with a non-coding region of the gene, such as an intron from the gene or 5' or 3' untranslated region (UTR).

Anti-sense strategies involve expression or transcription of an expression/transcription product capable of interfering with translation of mRNA transcribed from the target gene. This will normally be through the expression/transcription product hybridising to and forming a duplex with the target mRNA.

The expression/transcription product can be a relatively small molecule and still be capable of disrupting translation from the mRNA. However, the same result is achieved



by expressing the whole polynucleotide in an anti-sense orientation such that the RNA produced by transcription of the anti-sense oriented gene is complementary to all or part of the endogenous target mRNA.

- 5 Anti-sense strategies are described generally by Robinson-Benion *et al.* Methods in Enzymol 254, 363-375 (1995) and Kawasaki *et al.*, Artific. Organs 20, 836-845 (1996).

Genetic constructs designed for gene silencing may include an inverted repeat. An 'inverted repeat' is a sequence that is repeated where the second half of the repeat is in the complementary strand, e.g.,

5'-GATCTA.....TAGATC-3'

3'-CTAGAT.....ATCTAG-5'

The transcript formed may undergo complementary base pairing to form a hairpin structure provided there is a spacer of at least 3-5 bp between the repeated regions.

15

Another approach is to develop a small antisense RNA targeted to the transcript equivalent to an miRNA (Llave *et al.*, Science 297, 2053-2056 (2002) that could be used to target gene silencing.

- 20 The ribozyme approach to regulation of polypeptide expression involves inserting appropriate sequences or subsequences (eg. DNA or RNA) in ribozyme constructs (McIntyre Transgenic Res. 5 257-262 (1996)). Ribozymes are synthetic RNA molecules that comprise a hybridizing region complementary to two regions, each of which comprises at least 5 contiguous nucleotides of a mRNA molecule encoded by one of the inventive polynucleotides. Ribozymes possess highly specific endonuclease activity, which autocatalytically cleaves the mRNA.

25

Alternately, modulation may involve an increase in the expression and or activity of the polypeptide by over-expression of the polynucleotide, or by increasing the number of copies of the polynucleotide in the genome of the host.

30

To give effect to the above strategies, the invention also provides genetic constructs usually DNA constructs. The DNA constructs include the intended DNA (such as one or more copies of a polynucleotide sequence of the invention in a sense or anti-sense orientation or a polynucleotide encoding the appropriate ribozyme), a promoter sequence and a termination sequence (which control expression of the gene), operably linked to the DNA sequence to be transcribed. The promoter sequence is generally positioned at the 5' end of the DNA sequence to be transcribed, and is employed to initiate transcription of the DNA sequence. Promoter sequences are generally found in the 5' non-coding region of a gene but they may exist in introns (Luehrsen Mol. Gen. Genet 225, 81-93 (1991)) or in the coding region.

A variety of promoter sequences which may be usefully employed in the DNA constructs of the present invention are well known in the art. The promoter sequence, and also the termination sequence, may be endogenous to the target plant host or may be exogenous, provided the promoter and terminator are functional in the target host. For example, the promoter and termination sequences may be from other plant species, plant viruses, bacterial plasmids and the like. Preferably, promoter and termination sequences are those endogenously associated with the multifunctional germacrene-D synthase genes.

Factors influencing the choice of promoter include the desired tissue specificity of the construct, and the timing of transcription and translation. For example, constitutive promoters, such as the 35S Cauliflower Mosaic Virus (CaMV 35S) promoter, will affect the transcription in all parts of the plant. Use of a tissue specific promoter will result in production of the desired sense or antisense RNA only in the tissue of interest. With DNA constructs employing inducible promoter sequences, the rate of RNA polymerase binding and initiation can be modulated by external stimuli, such as light, heat, anaerobic stress, alteration in nutrient conditions and the like. Temporally regulated promoters can be employed to effect modulation of the rate of RNA polymerase binding and initiation at a specific time during development of a transformed cell. Preferably, the original promoters from the gene in question, or promoters from a specific tissue-targeted gene in the organism to be transformed are used. Other

examples of promoters which may be usefully employed in the present invention include, mannopine synthase (mas), octopine synthase (ocs) and those reviewed by Chua *et al.* Science 244, 174-181 (1989).

- 5 The termination sequence, which is located 3' to the DNA sequence to be transcribed, may come from the same gene as the promoter sequence or may be from a different gene. Many termination sequences known in the art may be usefully employed in the present invention, such as the 3' end of the *Agrobacterium tumefaciens* nopaline synthase gene. However, preferred termination sequences are those from the original  
10 gene or from the target species to be transformed.

The DNA constructs of the present invention may also contain a selection marker that is effective in cells, to allow for the detection of transformed cells containing the construct. Such markers, which are well known in the art, typically confer resistance to  
15 one or more toxins. One example of such a marker is the NPTII gene whose expression results in resistance to kanamycin or hygromycin, antibiotics which are usually toxic to plant cells at a moderate concentration. Alternatively, the presence of the desired construct in transformed cells can be determined by means of other techniques well known in the art, such as PCR or Southern blots.

20 Techniques for operatively linking the components of the inventive DNA constructs are well known in the art and include the use of synthetic linkers containing one or more restriction endonuclease sites. The DNA construct may be linked to a vector capable of replication in at least one host system, additional to the intended destination host system  
25 for example, *E. coli*, whereby after each manipulation the resulting construct can be sequenced and the correctness of the manipulation determined.

The DNA constructs of the present invention may be used to transform a variety of plants including agricultural, ornamental and horticultural plants. In a preferred  
30 embodiment, the DNA constructs are employed to transform kiwifruit, blueberry, apple, banana, tomato, cotton, rose, olive, potato, carnation, mango, papaya, freesia, orchids, lisianthus, gerbera, grape plants for altered flavour, fragrance, disease or insect

attraction characteristics. In addition the DNA constructs of the present invention may be used to transform medicinal plants such as *Solidago* species or plants used for biopharming such as tobacco. In a particularly preferred embodiment the plant is a carnation.

5

As discussed above, transformation of a plant with a DNA construct including an open reading frame comprising a polynucleotide sequence of the invention wherein the open reading frame is orientated in a sense direction can, in some cases, lead to a decrease in expression of the polypeptide by co-suppression. Transformation of the plant with a DNA construct comprising an open reading frame or a non-coding (untranslated) region of a gene in an anti-sense orientation will lead to a decrease in the expression of the polypeptide in the transformed plant.

10

It will also be appreciated that transformation of other non-plant hosts is feasible, including well known prokaryotic and eukaryotic cells such as bacteria (e.g. *E. coli*, *Agrobacterium*, *Lactobacillus*), fungi, insect, and animal cells is anticipated. This would enable production of recombinant polypeptides of the invention or variants thereof. The use of cell free systems (e.g. Roche Rapid Translation System) for production of recombinant proteins is also anticipated (Zubay Annu Rev Genet 7, 267-287 (1973)).

15

20

The polypeptides of the invention produced in any such hosts may be isolated and purified from same using well known techniques. The polypeptides may be used in cell-free systems for enzymic synthesis of sesquiterpenes for flavouring or scent uses, or for use in pheromone or antimicrobial agents. Such compounds could be germacrene-D and/or *delta*-cadinene and/or *gamma*-cadinene, and/or *alpha*-cubebene and/or *delta*-elemene and/or *alpha*-ylangene and/or *alpha*-copaene and/or *gamma*-elemene and/or *gamma*-muurolene and/or selinadiene and/or *alpha*-muurolene and/or germacrene B, and/or *delta*-elemene and/or elemol and/or *alpha*-cadinol.

25

Techniques for stably incorporating DNA constructs into the genome of target plants are well known in the art and include *Agrobacterium tumefaciens* mediated introduction, electroporation, protoplast fusion, injection into reproductive organs, injection into

30

immature embryos, high velocity projectile introduction, floral dipping and the like. The choice of technique will depend upon the target plant to be transformed.

5 Once the cells are transformed, cells having the DNA construct incorporated into their genome may be selected by means of a marker, such as the kanamycin resistance marker discussed above. Transgenic cells may then be cultured in an appropriate medium to regenerate whole plants, using techniques well known in the art. In the case of protoplasts, the cell wall is allowed to reform under appropriate osmotic conditions. In the case of seeds or embryos, an appropriate germination or callus initiation medium is employed. For explants, an appropriate regeneration medium is used.

15 In addition to methods described above, several methods are well known in the art for transferring DNA constructs into a wide variety of plant species, including gymnosperms angiosperms, monocots and dicots. The resulting transformed plants may be reproduced sexually or asexually, using methods well known in the art, to give successive generations of transgenic plants.

20 The nucleotide sequence information provided herein will also be useful in programs for identifying nucleic acid variants from, for example, other organisms or tissues, particularly plants, and for pre-selecting plants with mutations in multifunctional germacrene-D synthase genes or their equivalents which render those plants useful in an accelerated breeding program to produce plants in which the content of germacrene-D or *delta*-cadinene (and other cpds) and their derivatives is modulated. More particularly, the nucleotide sequence information provided herein may be used to design probes and primers for probing or amplification of multifunctional germacrene-D synthase encoding polynucleotides. An oligonucleotide for use in probing or PCR may be about 30 or fewer nucleotides in length. Generally, specific primers are upwards of 14 nucleotides in length. For optimum specificity and cost effectiveness, primers of 16-24 nucleotides in length are preferred. Those skilled in the art are well versed in the design of primers for use in processes such as PCR.

If required, probing can be done with entire restriction fragments of the gene disclosed herein. Naturally, sequences based upon Figure 3 (SEQ ID NO:1) or the complements thereof can be used. Such probes and primers also form aspects of the present invention.

5

Methods to find variants of the of polynucleotides of the invention from any species, using the sequence information provided by the invention, include but are not limited to, screening of cDNA libraries, RT-PCR, screening of genomic libraries and computer aided searching of EST, cDNA and genomic databases. Such methods are well known to those skilled in the art. Such variants are exemplified in Example 2.

10

The invention will now be illustrated with reference to the following non-limiting Examples.

## 15 EXAMPLES

### EXAMPLE 1

**Plant material and volatile collection:** *A. deliciosa* ([A. Chev] C.F. Liang et A.R. Ferguson var. *deliciosa* 'Hayward') flowers were taken at anthesis from vines grown in a HortResearch orchard at Te Puke New Zealand in 2003. Branches containing flowers were transported with stems in water, immediately after harvest. Whole flowers (at all stages of petal unfolding, and in good quality; 124.9g) were picked just under the receptacle and placed into a 250 mL Quickfit® Erlenmeyer flasks which was fitted with a headspace adaptor with an air inlet and outlet. Volatiles were collected in duplicate onto 100mg Chromosorb 105 adsorbent traps, which were fitted to the air outlet port of each adaptor. The volatiles were purged from the vessel onto the traps with clean air at 100 mL min<sup>-1</sup> for 60 minutes. The traps were stored at -15°C for one week, and then subjected to GC-FID/MS analysis.

20

25

Whole flowers of the same plants were also harvested at anthesis in 2000. 18.6g and 21g of flowers from a mixture of developmental stages were placed in 2 x Quickfit® Erlenmeyer flasks as above. Volatiles were collected onto a Chromosorb adsorbent trap

30

in duplicate, after the system had been equilibrated for 30 min at room temperature (ca. 23°C), by purging onto the trap with clean air at 25 mL min<sup>-1</sup> for 60 minutes. The trap was stored at -15°C for one week and then analysed by GC-FID/MS. The same petals (1.99g) were then extracted with pentane-ether (2ml) and stored at -20°C until analysed.

5 The pentane (GPR, BDH) had been purified by distillation and by passing through a column of activated alumina. Pentane extracts were reduced by rotary evaporation and were dried by passing through a small column of anhydrous MgSO<sub>4</sub> before the volume was reduced to ca. 50 µL under a gentle stream of nitrogen. The extract was analysed by GC-FID/MS.

10 Whole flowers of *A. deliciosa* var *deliciosa* genotype 32-10-07a were also collected in 2000. 3.5g of flowers from mixed developmental stages were placed in a Quickfit® Erlenmeyer flask as above. Volatiles were collected onto a Tenax TA adsorbent trap, after the system had been equilibrated for 30 min at room temperature (ca. 20°C), by  
15 purging onto the trap with clean air at 20 mL min<sup>-1</sup> for 100 minutes. The trap was stored at -15°C for two weeks and then analysed by GC-FID/MS.

**GC-MS Analysis of dynamic headspace samples and pentane extracts:** Volatile compounds were thermally desorbed from the headspace traps at 150 °C and were cryo-  
20 focussed at the beginning of the GC column (Young, 1981). The column outlet was split between the GC (Hewlett Packard 5890) FID detector (for quantitation) and VG-70SE (VG-Micromass, Manchester, U.K.) mass spectrometer (for component identification) with an electron impact ionisation of 70 eV. Separations were carried out in a 30 m x 0.32 mm i.d., 0.5 µm DBWax capillary column (J & W Scientific, Folsom, USA),  
25 starting at 30 °C for 6 min, increasing by 3 °C min<sup>-1</sup> to 102 °C, 5 °C min<sup>-1</sup> to 210 °C and held for 5 min. Injector temperature was 210 °C. The carrier gas was He at a flow rate of ~30 cm s<sup>-1</sup>, and the FID and mass spectrometer transfer line were at 210°C. Quantification of compounds was carried out using an average detector response based on methyl butanoate, ethyl butanoate, hexanol and methyl benzoate. Components were  
30 identified by comparison with spectra in the mass spectral database (NIST 1998, Wiley 7 and an in-house database), retention indices (in-house database) and in some cases direct GC-MS comparison with authentic standards. Separations for the earlier 2000

samples were carried out in the same manner as above except the programme started at 30°C for 6 min, increasing by 3°C min<sup>-1</sup> to 102°C, 5 °C min<sup>-1</sup> to 190°C and held for 5 min. Analysis of the pentane extract used a programme of 2 min at 40°C, 5 °C min<sup>-1</sup> to 200°C, 15 °C min<sup>-1</sup> to 240°C and held for 20 min. For genotype 32-10-07a: the GC-MS was programmed to start at 30 °C for 2 min, increasing by 3 °C min<sup>-1</sup> to 50 °C, 5 °C min<sup>-1</sup> to 130 °C, 10 °C min<sup>-1</sup> to 240 °C and held for 2 min. Injector temperature was 240 °C.

## RESULTS:

10 **Headspace analysis of volatiles emitted from kiwifruit flowers:** Germacrene D is a commonly found constituent of plants and exists as one of two isomers. (S)-(-)-germacrene D mainly occurs in higher plants, whereas (R)-(+)- germacrene D is predominantly found in lower plants such as liverworts (König *et al* Phytochem 43, 629-633 (1996)). In the headspace of the *A. deliciosa* flowers analysed, germacrene D  
15 of unknown enantiomeric composition was detected, at an average concentration of 0.68 ng/g of petal tissue. In the total ion chromatogram, TIC (see Figure 7a) the germacrene D is unresolved from the peak at 44.42 minutes but can be clearly identified by selecting ion 161 which is the base peak for Germacrene D (see Figure 7b) from the total ion chromatogram. Germacrene D was identified as occurring at 44.22. Other major  
20 sesquiterpenes also show fragment ions at 161 and there is another major sesquiterpene in *A. deliciosa* flowers. This was identified as *alpha*-farnesene.

A number of minor sesquiterpenes were found in the flowers, these were *delta*-elemene, *beta*-elemene, germacrene B and elemol. Tentative identifications were also given to  
25 some trace level sesquiterpenes found, these were *beta*-cubebene, calarene (aka *alpha*-gujunene) and *alpha*-humulene. Three trace level sesquiterpenes remained unidentified. Some monoterpenes were present; these were *beta*-myrcene, *alpha*-terpinene, limonene and 2 isomers of ocimene. An identical experiment carried out previously in 2000 showed *delta*-elemene, *beta*-elemene, germacrene B, *beta*-citronellol and one  
30 unidentified sesquiterpene present in the flowers in addition to the higher levels of germacrene D and *alpha*-farnesene. Pentane extracts of this material showed germacrene D (592 ng/g), *alpha*-farnesene (23669 ng/g) and elemol (3353 ng/g) to be



present. Both of these results confirmed Tenax trap extracts from another genotype of *A. deliciosa* ([A. chev] C.F. Liang et A.R. Ferguson var. *deliciosa*) flowers collected in a similar manner in 2000. These too showed Germacrene D and *alpha*-farnesene as the only major sesquiterpenes present.

5

## EXAMPLE 2

### EST library construction and analysis of gene and peptide characteristics including variants and phylogenetic analysis

- 10 **Isolation of mRNA and construction of cDNA library:** Total RNA was extracted from *A. deliciosa* petals by an adaptation of the method of Gomez and Gomez (Langenkamper, *et al.*, 36, 857-869 (1998)). mRNA was purified from the total RNA by oligo(dT)-cellulose chromatography (Pharmacia) and was used to construct a Lambda ZAP-CMV (Stratagene) cDNA library according to the manufacturer's instructions.
- 15 The cDNA-containing pBK-CMV plasmids were massed excised and used to transform *Escherichia coli* XL0LR (Stratagene). Plasmid was isolated from the XL0LR colonies and partially sequenced; the resulting cDNA sequences were then entered into the HortResearch EST database. All sequences on the database were compared to the NRDB90 database using the BLAST program (Altschul, Stephen F., Thomas L.
- 20 Madden, Alejandro A. Schaffer, Jinghui Zhang, Zheng Zhang, Webb Miller, and David J. Lipman (1997), "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs", *Nucleic Acids Res.* 25:3389-3402.). Putative terpene synthase cDNA sequences were identified by their similarity to known terpene synthases based on key protein motifs. A full-length sequence encoding a terpene
- 25 synthase (EST75565) was identified. EST75565 was fully sequenced to determine that a functional cDNA suitable for *in vitro* expression studies had been isolated SEQ ID NO: 1, Figure 3. Two other variants isolated from the *A. deliciosa* 'Hayward' petal cDNA library, ESTs 72838 and 80968 were fully sequenced also (SEQ ID NO:3 and SEQ ID NO:4, Figure 5). The encoded amino acid sequences are shown in Figure 4
- 30 (SEQ ID NO. 2) and Figure 6 (SEQ ID NO: 5 and SEQ ID NO: 6).

**Computational analysis of sequence information:** Computational analysis was performed using the European Molecular Biology Open Software Suite (EMBOSS). (Rice *et al*, 2000). The Transeq program was used to determine the amino acid sequence and the Pepstats program was used for the determination of molecular weight and isoelectric point. Sequence identity and similarity was calculated using the pair wise alignment program Needle, which uses the algorithm of Needleman and Wunsch (J. Mol. Biol. 48; 443-453 (1970)). The default parameters were used (Gap extension penalty: 0.5 for any sequence; Gap opening penalty: 10 for any sequence). Amino acid sequence alignments were performed using the program CLUSTALW (Thompson *et al*, 1994), and trimmed and shaded using the program GeneDoc (Nicholas and Nicholas, 1997).

## RESULTS:

**Sequence analysis of multifunctional germacrene D synthase:** Sequencing of EST75565 revealed an insert size of 1995 base pairs excluding the poly (A) tail. The cDNA sequence had an ORF encoding 565 amino acids with a putative start codon 82 bases downstream of the 5' end. The molecular mass of EST75565 was calculated to be 65 kDa. The predicted amino acid sequence of the multifunctional germacrene D synthase does not contain a chloroplast-signalling peptide sequence (Emanuelsson, Nielsen, Brunak and von Heijne, 2000). These signalling peptides are typical of monoterpene and diterpene synthases and its absence is suggestive of it being a sesquiterpene synthase. As has been found for most other terpene synthases the predicted amino acid sequence of the multifunctional germacrene D synthase also contains a DDX(X,D,E) motif (DDIYD) at amino acids 317 to 321 (Fig 4, SEQ ID NO:2). This motif is involved in the binding of divalent metal ions necessary for catalysis. Multifunctional germacrene D synthase was also shown to contain the angiosperm sesquiterpene consensus sequence GVIYXEP (GVYFEP) (Cai *et al* Phytochem 61, 523-529 (2002)), from amino acids 292 to 297.

Bohlmann, Meyer-Gauen and Croteau (Proc. Natl. Acad. Sci. USA 95, 4126-4133 (1998)) compared the amino acid sequences of 33 terpene synthases and showed that

there were seven absolutely conserved amino acid residues. Our multifunctional germacrene D synthase contains all seven of these absolutely conserved amino acids (Fig 4). They also found that six positions were absolutely conserved for aromatic amino acids and four positions were absolutely conserved for acidic amino acids. In our multifunctional germacrene D synthase, four of the six aromatic positions and three of the four acidic positions are conserved.

Several variants were also isolated. EST 80968 is a full length homologue of EST 75565 (SEQ ID NO: 4) with 95% identity at the amino acid level (SEQ ID NO:6) and 97% similarity. It is predicted to form a protein of the same size. It contains the same GVPYFEP motif from amino acids 292 to 297, and the DDIYD metal binding motif at amino acids 317 to 321. The ORF encoding 565 amino acids has a putative start codon 123 bases downstream of the 5' end. The EST is 1897 base pairs long. EST 72838 is a further full length homologue of EST 75565 with an exon deleted (SEQ ID NO:3).

Where there are corresponding regions the proteins are 98% identical and 99% similar (SEQ ID NO:5). It does not contain the GVPYFEP or DDIYD motifs as these are situated in the deleted exon. The 1817 base pair insert has a predicted ORF of 491 amino acids with a putative start codon 174 downstream from the ATG. Based on the untranslated regions the ESTs appear to be allelic variants of EST75565 (SEQ ID NO:1).

The predicted isoelectric point for multifunctional germacrene D synthase is 5.6 which is similar to the isoelectric point calculated for other sesquiterpene synthases. For example, two sesquiterpene synthases isolated from *Artemisia annua*, cASC34 and cASC125, have isoelectric points of 5.28 and 5.50, respectively (Van Geldre, *et al.*, Plant Sci. 158, 163-171 (2000)).

BLAST searches revealed that the predicted amino acid sequence of germacrene/cadinene synthase is most similar to the rose germacrene-D synthase (Guterman *et al.*, Plant Cell 14, 2325-2338 (2002)), having 59 % identity and 74 % similarity. Two sesquiterpene synthases from *Gossypium arboreum* (tree cotton), *delta*-cadinene synthase (Liang, Genbank, 1998), and *delta*-cadinene synthase isozyme A

(Chen *et al.*, J. Nat. Prod. 59 (10), 944-951 (1996)) with only two base pairs of difference, show the second highest similarity having 52% identity and 69% similarity at the amino acid level for both proteins. A sesquiterpene synthase from *Lycopersicon hirsutum* (tomato) of unknown function has the third highest similarity (van Der Hoeven *et al.*, Plant Cell 12 (11), 2283-2294 (2000)), with 49% identity and 69% similarity. Specific regions of amino acid identity between these synthases are generally short (usually between 5 and 10 amino acids). The most notable homologies between all four proteins is a region immediately upstream of and including the putative angiosperm sesquiterpene synthase consensus sequence GVFYFEP (Cai *et al* Phytochem 61, 523-529 2002). This includes a region of 13 or 14 identical amino acids for rose germacrene D synthase and the *Lycopersicon hirsutum* sesquiterpene synthase. The rose germacrene-D synthase also shares a region of 16 identical amino acids with EST 75565 from amino acids 314 to 329. Other regions of identity between these synthases include 4 amino acids from amino acid 167 to 170 and 4 amino acids from amino acid 265 to 268.

Homologues to EST 75565 were searched for among ESTs from other fruit species cDNA libraries. Two variants were found: EST 304951, a partially sequenced EST isolated from active *A. chinensis* 'Hort16A' meristems (Figure 5 and 6, SEQ NO: 7 and SEQ NO: 8) and EST 82293, a partially sequenced EST from blueberry (*Vaccinium corymbosum* 'Duke' fruit (Figures 5 and 6, SEQ ID NO: 9 and SEQ ID NO: 10)). Both sequence fragments start after the DDIYD motif at amino acid 375 (*A. chinensis*) and amino acid 395 (blueberry).

Phylogenetic analysis using the full length protein against known characterized and putative terpene synthases clustered multifunctional germacrene D synthase with rose germacrene D synthase, and *Gossypium* cadinene synthases in terpene synthase family a. The separation between rose and *A. deliciosa* germacrene D synthases and the cadinene synthases was well supported (95% of the time segregating with 1000 bootstraps in a neighbour joining unrooted tree), and the separation between the two germacrene D synthases was also well supported (100% of the time segregating with 1000 bootstraps).

**EXAMPLE 3.**

**Volatiles produced by bacterial cultures and purified protein Cloning into pGEM-T Easy and pET-30 vectors:** For functional expression, a 1735 bp cDNA fragment encoding EST75565, excluding the initiating methionine, was amplified by PCR using the following primers: forward primer: 5'- GAA TTC CAA CTA CCT TGT GCT CAA GC - 3' (SEQ ID NO:11); and reverse primer: 5' - CTC GAG CCT CCA CTT CAG TGT CTT G - 3' (SEQ ID NO: 12) (restriction sites underlined). PCR reactions were carried out in a total volume of 50 mL with the following reagents: 1x Expand High Fidelity PCR buffer (Boehringer), 0.2 mM dNTP's (Boehringer), 0.2 µM of each primer, and 1.75 units of Expand High Fidelity polymerase (Boehringer). PCR cycling was 94 °C (4 min); 25 cycles of 94 °C (30 s), 55 °C (30 s), 72 °C (2 min); and a final extension period at 72 °C (10 min). PCR products were purified using the QIAquick PCR clean up system (Qiagen) following the manufacture's specifications and cloned into pGEM-T Easy (Promega). A pGEM-T Easy clone harbouring a potential cDNA was fully sequenced to check for PCR errors, then excised using *Eco* RI and *Xho* I and subcloned into the expression vector pET-30a, yielding plasmid pET-30a75565. pET-30a75565 was resequenced at the 5' end to ensure the inserted cDNA was in frame, and then transformed into *E. coli* BL21-CodonPlus<sup>TM</sup>-RIL cells (Stratagene).

**Characterization of multifunctional germacrene D synthase from bacterial extracts and purified recombinant protein:** *E. coli* BL21-Plus<sup>TM</sup> -RIL cells harbouring pET-30a75565 were grown overnight at 37°C in Lauria-Bertani media supplemented with 30 µg ml<sup>-1</sup> kanamycin and 34 µg ml chloramphenicol<sup>-1</sup>. 5 mL aliquots of overnight culture were used to inoculate 4x 300 mL of fresh 2 x YT medium supplemented with 30 µg ml<sup>-1</sup> kanamycin and 34 µg ml<sup>-1</sup> chloramphenicol in 1 L baffled flasks. Cultures were grown at 37 °C with vigorous agitation to A<sub>600</sub> = 0.8, then removed to 4 °C to equilibrate to 16 °C before induction with 0.3 mM IPTG. Following induction, cultures were immediately transferred to a 16°C incubator and allowed to grow for a further 24 hours with continuous agitation and then harvested by centrifugation (2000 x g for 10min). Pelleted cells were resuspended in 20 mL binding buffer (5 mM imidazole, 0.5 mM NaCl, 10 mM DTT, 20 mM Tris-HCl (pH 7.9). Cells were disrupted with 2 times exposure to 12,700 psi in a French Pressure Cell Press (American Instrument Co. Inc,

Silver Spring, Maryland, USA) and then centrifuged at 10000 x g for 15 min. 5 mL of supernatant was transferred to a 50 mL test-tube and adjusted to 10 mM MgCl<sub>2</sub> and 20 µM MnCl<sub>2</sub>. FDP (or GDP) was added to a final concentration of 100 µM and the reaction mixture was incubated at 30 °C with shaking (110 rpm). Headspace volatiles were collected as below. The remainder of the extract (15 mL) was applied to a PD-10 gel filtration column (Amersham-Pharmacia Biotech) pre-equilibrated with binding buffer (DTT omitted). Eluent fractions were then pooled and purification of recombinant protein was carried out in a single step using immobilised metal affinity chromatography (IMAC) utilising the recombinant proteins hexa-histidine (His<sub>6</sub> or His<sub>6</sub> tag). The eluent was applied to a Hi-Trap Chelating HP column (Amersham-Pharmacia Biotech) charged with Ni<sup>2+</sup>. Non-bound proteins were removed and recombinant protein was eluted following the manufacturer's specifications. 5 mL samples of the eluted protein were transferred to 50 mL test tubes and adjusted to 100 µM FDP (or GDP), 10 mM MgCl<sub>2</sub>, 20 µM MnCl<sub>2</sub>. Optimum activity conditions are described in EXAMPLE 5. Headspace volatiles were collected as below. A 5 mL purified sample was also trapped without the addition of any precursors as a further control. Aliquots of the remaining protein fraction were stored at -80°C in 20% glycerol until required for further analysis.

**Electrophoresis and western analysis:** Whole culture, His-purified and non His-purified protein extracts were analysed by SDS-PAGE. Protein bands were separated on a polyacrylamide gel comprising a 4% stacking layer and a 10% separation layer. Protein bands were either visualised using modified Neuhoff stain or were transferred on to Immobilon-P PVDF membrane (Millipore) using a Trans-Blot semi-dry electrophoretic transfer cell (Bio Rad). Blotted proteins were incubated with Anti-His<sub>6</sub> monoclonal antibody (Roche) and then with Anti-Mouse IgG-AP (Stressgen) secondary antibody, and were detected using 1-STEP™ NBT/BCIP (Pierce) alkaline phosphatase detection reagent.

**Protein quantification:** The protein concentration of crude and partially purified protein extracts were determined using the Bio Rad Protein-Assay™ reagent according to the manufacturers specifications. The reactions were quantified at 595 nm with a

SPECTROmax PLUS spectrophotometer (Molecular Devices) using bovine serum albumin (BSA) as a standard.

#### Headspace trapping of purified protein

The headspace in the testubes was collected immediately after the addition of FDP (or GDP) and hexadecane (internal standard) using solid phase micro extraction (SPME). The SPME fibres (65 µm Carbowax®/DVB Supelco, Australia) were conditioned 30 min at 220 °C. The background was analysed for contamination using GC-FID (HP5890) prior to use. The headspace volatiles were collected for ~20 hours at 30°C with continuous agitation (110 rpm). Prior to analysis using a GC-FID/MS, the fibres were stored at ambient temperature in septum sealed glass vials.

#### GC-MS Analysis of SPME fibre samples

The volatiles were desorbed from the fibres in the GC injection port for 10 minutes at 220 °C for the Carbowax/DVB fibres. The GC system was equipped with a DB-Wax capillary column (J & W Scientific, Folsom, USA), 30 m x 0.25 mm i.d., with a 0.5 µm film thickness. The carrier gas was helium at a flow rate of 30 cm s<sup>-1</sup>. The GC oven was programmed to remain at 30°C for 6 min, then to increase by 3 °C min<sup>-1</sup> to 102 °C, followed by an increase of 5 °C min<sup>-1</sup> to 210 °C, which was maintained for 11 min. The mass spectrometer operated in electron impact ionisation (EI-MS) mode at 70 eV with a scan range of 30 – 320 amu. Peak identification was carried out by comparison of sample spectra with those from NIST98, Wiley 7, and our own mass spectra libraries and was confirmed by retention indices of authentic standards and literature values (Davies 1998). Semi-quantitative data was obtained by measuring sample peak area relative to hexadecane internal standard.

#### Dynamic headspace trapping and pentane extraction and identification of volatiles produced by the recombinant multifunctional germacrene D synthase protein.

Dynamic headspace methods were also used to collect the headspace volatiles from the purified extracts on to 100mg Chromosorb 105 adsorbent traps. The protein was placed into 50 mL Quickfit® test tubes, which were fitted with a headspace adaptor with an air inlet and outlet. Volatiles were collected onto the Chromosorb 105 adsorbent traps, which were fitted to the air outlet port of each adaptor. The volatiles were purged from

the vessel onto the traps with clean air at 50 mL min<sup>-1</sup> for 18 hours. The traps were stored at -15°C for up to one week, and then subjected to GC-FID/MS analysis. Finally, pentane extracts were taken from the purified extracts. An aliquot (2.5ml) of pentane was added to the purified extracts and was left to extract for 18 hours. The pentane was removed and then concentrated with a stream of N<sub>2</sub> gas. This was then injected onto the GC-MS at 150 °C.

## RESULTS

**Western analysis:** Western analysis confirmed the presence of a soluble expression product that was slightly larger (ca. 70 kDa) than the predicted 65kDa for the native protein. This size difference can be attributed to a His<sub>6</sub> tag being present in the recombinant EST75565 expression product. This product was detected in both the bacterial and partially purified recombinant protein extracts. No equivalent bands were detected in any of the vector only control extracts.

### Characterisation of multifunctional germacrene D synthase:

When the purified protein sample which contained pET30/75565 and the FDP precursor was analysed for presence of compounds in the volatile headspace, a significant new peak was found, when compared to the control of empty vector protein and FDP.

Library identification of this peak showed that it was germacrene D (see Figure 8). When trapping using the Carbowax®/DVB fibres Germacrene-D (83%), a lesser amount of *gamma*-elemene (1.3%), *delta*-elemene (7%), germacrene-B (4.3%), elemol (4.2%) and *delta*-cadinene (0.5%) and *gamma*-muurolene (0.3%) were identified (see Figure 8).

The profile given by the dynamic headspace trapping method was quite similar to that of the Carbowax®/DVB SPME fibres, in that Germacrene D was the main product (54%) with smaller amounts of *delta*-elemene (15.5%), *gamma*-elemene (2.9%), Germacrene B (1.4%), *delta*-cadinene (3.9%) and elemol (16.6%) (Figure 9). The levels of *delta*-elemene, *gamma*-elemene, *delta*-cadinene, *gamma*-muurolene and elemol were higher using this method but germacrene D was much lower than found with the SPME fibre. *Gamma*-muurolene (3.8%) was at a much higher percentage in



this sample and *gamma*-cadinene (1.8%) was also identified; it could not be identified in the Carbowax®/DVB SPME fibres. This difference in ratios is likely to be due to the chemical properties and specificity of the trapping matrix

- 5 The decrease of germacrene D and concomitant increase in *delta*-elemene, *gamma*-elemene, *delta*-cadinene, *gamma*-muurolene and elemol may suggest that these compounds are rearrangements of germacrene D, in particular thermal, photochemical or acid catalysed. It is well established that thermal rearrangement of germacrene compounds leads to the corresponding elemene compounds (germacrene A rearranges
- 10 to *beta*-elemene, germacrene B rearranges to *gamma*-elemene and germacrene C rearranges to *delta*-elemene), however, germacrene D does not thermally rearrange to an elemene structure. Germacrene A and C are very thermally labile, hence require low temperature GC-MS analysis (<100 °C) (Feger *et al.*, Flavour and Fragrance Journal 15: 170-173 (2000)) to ascertain whether these products are due to rearrangement of the
- 15 corresponding germacrenes. Although germacrene D is stable up to 180 °C, germacrene D can thermally rearrange to *beta*-ylangene, *beta*-copaene, *epsilon*-muurolene and *epsilon*-amorphene as major products (Bülow and König, Phytochem 55: 141-168 (2000)). The comparison of the products on the Chromosorb cartridge (which was desorbed at 150 °C) and the Carbowax®/DVB SPME fibre (which was
- 20 desorbed at 220 °C) does not reveal any differences which can be attributed to thermal rearrangement of germacrene D. In terms of the acid catalysed rearrangement, germacrene D completely rearranges to different sesquiterpenes by contact with weakly acidic media over a period of several days (Yoshihara *et al.*, Tetrahedron Letts. 27: 2263-2264 (1969)), the main products of acid catalysed rearrangements are cadinene,
- 25 muurolene and amorphene structures. Cadinenes are the predominant products due to a greater thermodynamic favorability of forming the cadinyl cation compared to the formation of the muurolenyl and amorphenyl cations. Although the acidity of the 2 matrices may have caused some rearrangement to the cadinene type structures, the extractions were introduced to the MS within a matter of hours, and germacrene D very
- 30 often occurs with *delta*-cadinene in plants (Yoshihara and Hirose, Bull. Chem. Soc. Jpn, 51: 3395-3396 (1978)).

More light can be shed on the possibility of rearrangement of the products using pentane extraction. Pentane extraction does not subject the products to different pHs and can be injected into the GC-MS at specific temperatures such that potential for thermal rearrangement can be assessed.

Table 1 Thermal rearrangements and peak areas for pentane extracts from purified protein

Compound	Injection Port Temperature (°C)		
	125	150	210
	Peak Area		
Germacrene D	$4.22 \times 10^6$	$4.90 \times 10^6$	$3.91 \times 10^6$
Germacrene B	$7.57 \times 10^5$	$8.58 \times 10^5$	$5.87 \times 10^5$
Elemol	$ca. 9 \times 10^4$	$ca. 1.4 \times 10^5$	$5.75 \times 10^6$
$\delta$ -Elemene	nd	nd	$9.05 \times 10^5$

<sup>nd</sup> compound not detected. The *ca.* values were for small elemol peaks that were not fully resolved and thus are only approximations.

Table 2 Thermal rearrangements and peak areas for pentane extracts from *Actinidia* flowers.

Compound	Injection Port Temperature (°C)			
	100	125	150	210
	Peak Area			
Germacrene D	$2.85 \times 10^7$	$3.78 \times 10^7$	$3.87 \times 10^7$	$2.42 \times 10^7$
Germacrene B	$5.0 \times 10^6$	$7.40 \times 10^6$	$1.0 \times 10^7$	$3.92 \times 10^6$
Elemol	nd	nd	$ca. 8.4 \times 10^6$	$3.78 \times 10^7$
$\delta$ -Elemene	nd	nd	$ca. 5.7 \times 10^5$	$4.14 \times 10^6$

<sup>nd</sup> compound not detected. The *ca.* values were for small peaks that were not fully resolved and thus are only approximations.

In an experiment to test for thermal rearrangement, injections of pentane extract from purified protein were made at temperatures of 125, 150 and 210°C; or pentane extract from *Actinidia* flowers were made at temperatures of 100, 125, 150 and 210°C. This demonstrated that while germacrene D remained at similar levels, elemol and *delta*-elemene increased in amount from non detectable (*delta*-elemene) or low levels (elemol) to significant quantities (Tables 1 and 2). Germacrene B modulated slightly in response to temperature shifts although no *gamma*-elemene was found as might be predicted. In addition the predicted precursor for *delta* elemene – germacrene C (Feger

*et al.*, Flavour and Fragrance Journal 15: 170-173 (2000)) was not detected at any time. Germacrene A was also not detected at any time. This suggests that these elemene-based compounds may be present in our experiments due to thermal rearrangements, although the source of rearrangement is unclear. Elemol may arise from further rearrangements of *delta*-elemene or from other elemene sources or from hedycaryol (Chamblee *et al.*, J. Essent. Oil Res. 9: 127-132 (1997)). No GC-MS traces in any experiments indicated evidence for germacrene-C or for hedycaryol. Hence although the multifunctional germacrene D synthase may also produce germacrene C we have no evidence to support this other than the appearance of the elemene-based compounds at higher temperatures.

#### EXAMPLE 4

##### Effect Of Trapping Method On Products Detected.

**Expression and characterization of multifunctional germacrene D synthase from bacterial cultures:** *E. coli* BL21-Plus<sup>TM</sup>-RIL cells harbouring pET30a75565, or empty pET30a vector as a control, were grown overnight at 37 °C in Luria-Bertani media supplemented with 30 µg/mL kanamycin and 50 µg/mL chloramphenicol. A 500 mL aliquot of overnight culture was used to inoculate 50 mL of fresh 2 x YT medium supplemented with 30 µg/mL kanamycin and 50 µg/mL chloramphenicol. The culture was grown at 37 °C with vigorous agitation to A<sub>600</sub> = 0.6 before induction with 0.3 mM isopropyl-β-D-thiogalactopyranoside (IPTG) and simultaneous addition of both farnesyl diphosphate (FDP) (100 µM) and hexadecane (required as an internal standard) or geranyl diphosphate (GDP) and hexadecane. The culture was immediately transferred to an incubator set at 16 °C, 30 °C or 37 °C depending on the experiment

25

The headspace in the vessels above the bacterial cultures was collected immediately after the addition of FDP (or GDP) and hexadecane (the internal standard) using solid phase micro extraction (SPME). The SPME fibres (65 µm PDMS/DVB, Supelco, Australia) were conditioned for 45 min at 260 °C. The background was analysed for contamination using GC-FID (HP5890) prior to use. The headspace volatiles were collected for ~20 hours at 30°C with continuous agitation (110 rpm). Prior to analysis

30

using a GC-FID/MS, the fibres were stored at ambient temperature in septum sealed glass vials. The PDMS/DVB fibres were desorbed in the injector at 260 °C

As trapping whole cultures with PDMS/DVB fibres demonstrated the presence of compounds that had not been detected using the Carbowax®/DVB SPME fibres, other fibres were compared. The headspace of purified protein was also trapped using 85 µm Polyacrylate SPME and PDMS/DVB SPME fibres to allow comparison with the Carbowax®/DVB SPME fibres (Supelco, Australia). The polyacrylate SPME fibres were conditioned for 120 min at 300 °C and were desorbed in the injector at 300 °C.

## RESULTS

### *Bacterial cultures:*

Controls comprising *E. coli* BL21 cells transformed with pET-30a lacking the multifunctional germacrene-D synthase cDNA insert produced none of the compounds indicated by trapping of the purified protein with Carbowax®/DVB fibres (Figure 10a). Trapping the whole cultures containing the multifunctional germacrene-D synthase cDNA insert using the PDMS/DVB fibres gave rise to different compounds and different ratios compared to those detected from the purified recombinant protein using Carbowax®/DVB fibres (compare Fig 10b with Fig 8). In particular germacrene D was present in a lower ratio and *gamma*-muurolene appeared at a much higher level.

In this experiment the production of germacrene-D was the largest (37.6%). *Delta*-elemene (12.5%), a lesser amount of cadinene, primarily *delta* isomer (10.9%) but also the *gamma*- isomer (6.3%), germacrene B (9.1%), *gamma*-muurolene (5.7%), and smaller quantities (less than 5%) of a range of sesquiterpenes: *gamma*-elemene, selina-3,7(11)-diene, *alpha*-humulene, *alpha*-cadinene, *alpha*-muurolene, *alpha*- amorphene, *alpha*-ylangene, *alpha*-cubebene, elemol and 3 unknowns were also produced.

In the empty vector control, there were some sesquiterpenes produced, these were *cis-alpha*- and *beta*- bisabolene. There was no evidence of germacrene D or any of the products listed above. The internal standard indicated even efficiency of trapping for

both the pet30/75565 + FDP and pet30 control only + FDP samples. In addition, for the sample pet30/75565 with no added precursor, no germacrene D or any other sesquiterpenes were found.

- 5 Addition of geranyl diphosphate (GDP) to bacterial cultures did not result in the production of any monoterpenes. Germacrene-D was still produced, although at significantly lower levels to that exhibited with FDP addition. This is due presumably to the presence of bacterial FDP.

10 *Purified protein:*

Compounds released from incubation of purified protein with FDP were similar to those in whole cultures, although there were less background (bacterial) compounds present. In contrast to the addition of GDP to bacterial cultures, purified protein did not produce any germacrene D. This reinforces the theory that the germacrene D found in the  
15 pet30/75565 + GPP bacterial culture was only due to the presence of endogenous FDP in the bacteria.

**Effect of trapping matrix on sesquiterpene composition:** The enzyme produced different products depending on how the volatile products were extracted and analysed.  
20 When trapping with the PDMS/DVB fibre a range of sesquiterpene products were identified (Figure 11).

Of the 42 sesquiterpenes or sesquiterpene alcohols present, 84.7% of the products could be identified. The major products were *gamma*-muurolene (14.8%), *delta*-cadinene (14.2%), eudesma-3,7(11)-diene (aka selina-3,7(11)-diene (7.3%)), *gamma*-cadinene (7.3%), *alpha*-muurolene (5.9%), *gamma*-elemene (4.9%), germacrene-D (4.6%).  
25 Other compounds which were identified as products were *delta*-elemene, isomers of cadinol (including *alpha*-, *delta*- and *tau*-), *alpha*-cadinene, *alpha*-humulene, *alpha*-selinene, *alpha*-copaene, *alpha*-cubebene, elemol, germacrene-B, *alpha*-amorphene, -  
30 *alpha*-ylangene, *beta*-elemene, *beta*-cubebene, isomers of eudesmol and muurolol. Tentative identifications based on structure were given to epibicyclosesquiphellandrene and isodedene. Bicyclosesquiphellandrene is a known compound from acid catalysed

rearrangement of *delta*-cadinene or through the muurolenyl cation from germacrene D (Bulow and Konig, Phytochem 55: 141-168 (2000)). 18 compounds remained unidentified, only 2 of these compounds were in amounts over 1%, but these could not be resolved from each other hence no assignment could be made. Two of these eighteen  
5 compounds were sesquiterpene alcohols.

The polyacrylate fibre was found to be least efficient at extracting the sesquiterpenes, and showed low profiles of only germacrene D. The Carbowax®/DVB fibre showed germacrene D as the main product with smaller and similar amounts of *delta*-elemene,  
10 *gamma*-elemene, small amounts of *beta*-elemene, germacrene B, *delta*-cadinene and *gamma*-cadinene and elemol. Finally the PDMS/DVB fibre showed a wide range of products, with *gamma*-muurolene as the major product, followed by *delta*-cadinene, eudesma-3,7(11)-diene, *gamma*-cadinene, *alpha*-muurolene, *gamma*-elemene and germacrene-D. There were also a host of other sesquiterpenes which could potentially  
15 be attributed to acid catalysed rearrangements of Germacrene D, which may also account for the decrease in intensity of the Germacrene D on this fibre. Elemol was also detected but at very low amounts.

The differences in the SPME coatings relates to the acidity, and these results illustrate  
20 that germacrene D is capable of rearranging under certain conditions (Yoshihara et al., Tetrahed Lett 27: 2263-2364 (1969)), acid catalysis being an important rearrangement mechanism.

However, headspace trapping is a sensitive method for collecting the volatile products of enzymes which allows for preconcentration on the sampling medium and comparison  
25 of the different products obtainable with different extraction methods shows the scope of usability for this enzyme, in that the products obtained can be manipulated by the extraction method.

**Rearrangement with time of trapping:** The time courses (Table 3 and Figures 12 and  
30 13) show that over the period of time on PDMS/DVB fibres, the ratios of germacrene D become less compared to that of the *gamma*-muurolene and the 2 cadinene isomers, but *gamma*-muurolene and *delta*-cadinene maintain the same ratio relative to each other.

*Gamma*-muurolene is higher than *delta*-cadinene. This is not as predicted by Bulow and Konig (Phytochem 55: 141-168 (2000)) where *delta*-cadinene is 2-fold more prominent than *gamma*-muurolene in a silica catalysed rearrangement. In addition, elemol is not detectable with 4 hour trapping but is present after 12 hours and 24 hours.

5 This may be due to lack of equilibrium or slow kinetics.

Table 3. Relative quantities (%) of sesquiterpenes trapped onto PDMS/PDV fibres with time of trapping. Each trapping sampled for 30 minutes at 0, 1 and 2 hours after addition of FDP to the purified enzyme.

10	Compound	0	1	2
	<i>delta</i> -elemene	0	2.7	4.0
	<i>gamma</i> -elemene	0	5.6	5.9
	<i>gamma</i> -muurolene	15	20.2	20.0
	germacrene D	9.4	5.0	4.4
	unresolved	18.5	9.0	8.7
15	<i>alpha</i> -muurolene	0	6.9	8.3
	<i>alpha</i> -selinene	0	6.0	4.5
	<i>delta</i> -cadinene	12.6	15.8	16.6
	<i>gamma</i> -cadinene	14.5	13.2	12.3
	selinadiene	15.4	7.7	7.6
	<i>alpha</i> -cadinene	0	3.4	3.9
	germacrene B	14.6	4.5	3.8

20

#### EXAMPLE 5

**Optimisation of large-scale production of protein and protein kinetics:** *E. coli* BL21-Plus<sup>TM</sup> -RIL cells harbouring pET-30a75565 were grown overnight at 37°C in Lauria-Bertani media supplemented with 30 µg ml<sup>-1</sup> kanamycin and 34 µg ml chloramphenicol<sup>-1</sup>. 5 mL aliquots of overnight culture were used to inoculate 4x 300 mL of fresh 2 x YT medium supplemented with 30 µg ml<sup>-1</sup> kanamycin and 34 µg ml<sup>-1</sup> chloramphenicol in 1 L baffled flasks. Cultures were grown at 37 °C with vigorous agitation to A<sub>600</sub> = 0.8, then removed to 4 °C to equilibrate to 16 °C before induction with 0.3 mM IPTG. Induced cultures were then incubated at 16 °C and 220 rpm for a further 72 hours. Cells were pelleted by centrifugation (2500 x g; 10 min) and stored overnight at -20 °C. The following day cell pellets were resuspended in 20 mL His6 binding buffer and cells were disrupted by 2 passes through an EmulsiFlex®- C15 high-

pressure homogeniser (Avestin) with a pressure setting between 15000-20000 psi. Cell debris was pelleted by centrifugation 2x at 10000 x g for 15 min; 4 °C (Sorval SS34 rotor). The supernatant was filtered through a 0.45 µm filter (Amicon). Filtered extract was desalted and passed over a Nickel affinity column. Concentration of the recombinant synthase in the purified protein extracts was estimated from SDS-PAGE gels by comparing against a BenchMark™ Protein Ladder (Invitrogen). Purified extracts were adjusted to a workable concentration with His6 elution buffer (without Imidazole) containing 10 % glycerol and 1mM DTT. The extract was then separated into 100 µl aliquots and stored at -80 °C until required. As a control for future analysis, BL21-Plus™ -RIL cells harbouring pET-30a only were also induced and purified as above.

**Protein oligomerisation:** 1.5 mL of His6 purified protein extract (containing ~ 60 µg of the recombinant synthase purified protein) was loaded onto a 600 x 16 mm S200 Sephacryl column (Pharmacia) at a flow rate of 1 mL min<sup>-1</sup>. The column was pre-equilibrated and eluted with 50 mM Bis-tris propane buffer (pH7.5) containing 10 % glycerol and 150 mM KCl. Fractions corresponding to protein peak of interest were pooled and assayed after adjusting to 10 mM MgCl<sub>2</sub> and adding 20µM <sup>3</sup>H-FDP for activity. Molecular mass of active fractions was calculated based on comparison to standards of known molecular mass.

**Kinetic Studies:** For kinetic studies, Germacrene D synthase active protein that had been induced in culture and purified as described for protein optimisation was added to a minimal assay buffer containing 50 mM Bis-Tris- Propane (pH 7.5), 10 % (v/v) glycerol, 1 mM DTT and 0.1 % (v/v) Tween - 20. Tritiated FDP was added variously depending on the experiment. One mL assays containing 100 -150 ng of the recombinant synthase were overlaid with 0.6 mL pentane and incubated in 1.5 mL microfuge tubes for 2 hours at 30 °C and 150 rpm. All assays were performed in triplicate. Following incubation assays were immediately placed on ice and a 200 µL aliquot of the pentane layer removed for analysis. The aliquots were added to 1.5 mL microfuge tubes containing 0.7 mL Organic Counting Scintillant (OCS) (Amersham)



and vortexed briefly. Scintillation analysis was performed using a Wallac 1409 Liquid Scintillation Counter ( $^3\text{H}$  efficiency  $\approx 70\%$ ).

- Kinetic studies with  $^3\text{H}$ -FDP (10.06 Mbq/ mL) as substrate (concentration range 1  $\mu\text{M}$  to 100  $\mu\text{M}$  with saturating  $\text{Mg}^{2+}$  were carried out to determine  $K_m$  for FDP. Kinetic constants for  $\text{Mg}^{2+}$ , and  $\text{Mn}^{2+}$  at 10  $\mu\text{M}$   $^3\text{H}$ -FDP (assay range 50  $\mu\text{M}$  to 10 mM and 25  $\mu\text{M}$  to 5 mM of the chloride salts respectively) were determined. The effect on enzyme activity of metal co-factors with and without salts was also tested.  $\text{Mg}^{2+}$  and  $\text{Mn}^{2+}$  were added in the presence and absence of 50 mM KCL and 50 mM NaCl in all possible combinations. Controls included incubation without enzyme, with enzyme but without metal ion cofactors and with enzyme and cofactors in the presence of 10 mM EDTA. A further control comprising His6 purified pET-30a extract was also assayed to determine potential utilisation of FDP by co-purifying bacterial proteins.
- For determination of the enzyme pH optimum, assays were carried in a tri-buffer system containing 51 mM diethanolamine, 100 mM MES, and 51 mM N-ethylmorpholine at pH values between 5.7 and 9.5 with 10 mM  $\text{MgCl}_2$ , 10  $\mu\text{M}$  FDP, 10 % (v/v) glycerol, 1 mM DTT and 0.1 % (v/v) Tween - 20. Optimal temperatures for enzyme activity in the range 16  $^\circ\text{C}$  – 50  $^\circ\text{C}$  were also determined using the standard assay buffer and 10 mM  $\text{MgCl}_2$ , and 10  $\mu\text{M}$  FDP. Kinetic constants were determined from the DPM data by nonlinear regression using the Origin50 graphics package. Data presented represents the means of three determinations with standard errors within  $\pm 10\%$  and with background DPM calculated from controls subtracted.
- All experiments were carried out at least twice.

Because it was apparent that the matrix of the fibres could affect the composition of the sesquiterpenes trapped, we also checked the production of individual sesquiterpenes produced by purified protein in response to different pH conditions in the enzyme assay medium.

## RESULTS:

***K<sub>m</sub>*** The *K<sub>m</sub>* for FDP was in the 1 - 2  $\mu$ M range with saturating concentrations at  $\sim$  7.5  $\mu$ M (Figure 14A). The recombinant synthase exhibited a low *K<sub>i</sub>* with  $\sim$  50 % inhibition at 1 mM substrate concentrations. *K<sub>m</sub>* for the cofactor  $Mg^{2+}$  was in the 0.6 to 1.2 mM range (Figure 14B) and the *K<sub>m</sub>* for  $Mn^{2+}$ , although difficult to determine with a high degree of accuracy given very low synthase activity in the presence of  $Mn^{2+}$ , was in the 50 to 300  $\mu$ M range (Figure 14C). No significant inhibition was observed in the presence of high concentrations ( $>$  5 mM and 10 mM) of the respective  $Mg^{2+}$  and  $Mn^{2+}$  chloride salts. *K<sub>m</sub>*s in the literature range from 0.4 -4.5  $\mu$ M for FDP, 70 -150  $\mu$ M for Mg, 7-30  $\mu$ M for Mn for other sesquiterpene synthases (e.g. Cai et al Phytochemistry 61, 523-529 (2002), Steele et al J. Biol. Chem. 273, 2078-2089(1998)).

**pH** Germacrene D synthase showed a pH optimum of 7.5. pH values of 0.5 pH units lower and 1 pH unit higher than 7.5 resulted in 47 % and 59 % respective decreases in enzyme activity Figure 14D. This pH is within the range (between pH 7 and 9) reported for other characterised sesquiterpene synthases in the literature (Cai et al Phytochemistry 61, 523-529 (2002); Steele et al J. Biol. Chem. 273, 2078-2089 (1998))

**Temperature** Maximum synthase activity occurred at 30  $^{\circ}$ C, with a significant reduction in activity observed above 42  $^{\circ}$ C (Figure 15). Little or no activity was observed at 50  $^{\circ}$ C. Significant activity was observed at low temperatures however with the synthase exhibiting  $\sim$  56 % of the optimal temperature activity at 16  $^{\circ}$ C. The optimal temperature is similar to that reported for other sesquiterpene synthases.

**Other metal ion effects.** Germacrene D synthase has a requirement for  $Mg^{2+}$  as a cofactor for activity and inhibition by  $Mn^{2+}$  ions. Mn can act as a cofactor when alone, but inhibits Mg requiring cofactor activity. Addition of  $K^{+}$  or  $Na^{+}$  ions had little or no effect on activity, and the recombinant enzyme showed a strong preference for  $Mg^{2+}$  over  $Mn^{2+}$  (Vrel  $Mn^{2+}$   $\sim$  27 %). (Fig 16). The absence of DTT was also shown to significantly decrease activity (Vrel in the absence of DTT  $\sim$  53 % under optimal conditions. The presence of Mn in the buffer resulted in inhibition of activity even if  $Mg^{2+}$  was present.

**pH effect on ratios of sesquiterpene products:** Germacrene D was the major product when the volatiles produced by the purified enzyme were trapped at 3 different pHs onto Carbowax®/DVB fibres. Germacrene D was the major product at all 3 pH levels (Figure 17) and was present in similar percentages (Table 4).

The headspace profiles show that there is clearly less product at pH 8.5 and these results agree with the kinetic studies which suggest that the optimum pH is at pH7.5. However the other products, *delta*-elemene, *beta*-elemene, *gamma*-elemene, Germacrene B, *delta*-cadinene and elemol, varied in relative contribution with pH. The ratios of these products are given in Table 4

Table 4 Ratios of products at 3 different pHs trapped on Carbowax®/DVB SPME fibres after 21 hours

Compound	% at pH 6.5	% at pH 7.5	% at pH 8.5
<i>delta</i> -elemene	10.4	13.4	14.6
<i>beta</i> -elemene	10.6	6.3	1.7
<i>gamma</i> -elemene	5.9	6.8	9.0
germacrene D	64.6	64.8	64.9
germacrene B	2.7	2.3	4.8
<i>delta</i> -cadinene	nd	1.2	nd
elemol	5.9	5.2	4.9

#### EXAMPLE 6

##### Copy number and expression in *Actinidia*:

DNA was extracted from leaves of four *Actinidia* species (*Actinidia deliciosa* 'Hayward', *Actinidia chinensis* 'Hort16A', *Actinidia arguta* 'Hortgem Tahi' and *Actinidia* CK51\_05) by a modification of the plant DNA extraction method of Murray and Thompson Nucleic Acids Res. 8: 4321-5 (1980). The leaf material (up to 1g) was ground in liquid nitrogen and the powder was added to 20 ml buffer (2% cetyltrimethylammonium bromide, 100mM Tris pH 8.0, 20 mM EDTA, 1.4 M NaCl) that had been heated to 55°C. The extract was placed in a 55°C waterbath, 400 µl β-mercaptoethanol was added and the extract was gently swirled until the temperature of

the waterbath reached 50°C. The extract was then left at room temperature for 15 minutes, after which 20 ml chloroform:isoamyl alcohol (24:1) was added and the extract gently swirled until well-mixed. The extract was then centrifuged at 8000 rpm for 20 minutes at 20°C. The top layer was removed and an equal volume of isopropanol was added. After inverting gently until mixed the DNA was spooled out. RNase (Roche) (1 µl of 10 mg/ml) was added to remove RNA.

**Southern analysis:** Genomic DNA that had been extracted from leaves of four *Actinidia* species was put through the Wizard DNA Clean-Up system (Promega) and then digested overnight with *Eco*R1 in a total volume of 50µl. The digested DNA was then electrophoresed through 0.5% agarose, visualised with ethidium bromide, hydrolysed in 0.25M HCl and washed in water before transfer to Nytran-Plus (Schleicher & Schuell) membrane in 0.4M NaOH overnight. The membrane was then neutralised in 0.5M Tris and prehybridised in Washing and Pre-Hyb Solution (MRC) for two hours. Hybridisation was performed in 10 ml of High Efficiency Hybridization System (MRC) using as a probe a <sup>32</sup>P-labelled 350 base pair PCR fragment that was complementary to a region near the 5' end of the coding sequence of germacrene D synthase. The probe (40 ng) was labelled with <sup>32</sup>P dCTP using the *rediprime*<sup>TM</sup>II (Amersham Pharmacia) random labelling system, following the manufacturer's directions. The labelled probe was denatured in 0.1 M NaOH for 30 minutes before hybridisation overnight. The membrane was washed in Washing and Pre-Hyb Solution (MRC) according to the manufacturer's recommendations. Hybridisation signals were visualised by scanning on a Storm 840 phospho-imaging system (Molecular Dynamics) and analysed using ImageQuant software.

25

**RT-PCR amplification of multifunctional germacrene D synthase in plants:** RT-PCR (Platinum<sup>®</sup> Quantitative RT-PCR Thermoscript One-Step System, Invitrogen) amplifications were performed according to the manufacturer's recommendations on total RNA extracted from different tissues of *Actinidia deliciosa* and from *Actinidia chinensis* fruit and from the leaves of transgenic *Arabidopsis thaliana* transformed with EST 75565. cDNA synthesis was at 60°C for 30 minutes, followed by denaturation at

30

96°C for 5 minutes, then 40 cycles of amplification involving denaturation at 96°C for 30 seconds, annealing at 55°C for 40 seconds and extension at 72°C for 60 seconds. For the final cycle extension at 72°C was continued for a further 5 minutes. Prior to RT-PCR the total RNA was treated with DNase I (Life Technologies) for 10 minutes at room temperature. Concurrently with RT-PCR amplification, PCR amplification was also performed on the DNase I-treated total RNA to check for genomic DNA contamination. The PCR primers were as follows: 75565GSP1 CAATTGAGAGGTGGGAGATC (SEQ ID NO: 13) and 75565GSP2 GTTGGAAACATATCCTTGGTG (SEQ ID NO: 14) (Primer set 1); 75565NF1 TAGGCGTGTCTTACCATTTT (SEQ ID NO: 15) and 75565NR1 GCATGAATGATTGTTCCTT (SEQ ID NO: 16) (Primer set 2). PCR primer 75565GSP1, which is specific for EST 75565, was designed to amplify over an intron splice site to prohibit genomic DNA amplification. The resulting amplification products (5 µl) were analysed by electrophoresis through 1% agarose (AppliChem), followed by staining with ethidium bromide and visualisation on an ultraviolet transilluminator (UVP) attached to a camera (UV tec, Total Lab Systems Ltd).

## RESULTS:

Southern analysis revealed that the multifunctional germacrene D synthase was present at low copy number in all four genotypes tested. A homologous gene was clearly also present in *A. arguta* although EST sequencing had not revealed its presence.

The multifunctional germacrene D synthase was highly expressed in *A. deliciosa* 'Hayward' petals (Figure 18). Lower concentrations of transcript were detected by northern analysis in stamens and styles. The more sensitive PCR technique indicated low levels of transcript in ovaries and mature leaves of 'Hayward' with one set of primers. A second set of primers however detected transcripts in expanded leaves and young fruitlets (to 5 weeks after anthesis) in 'Hayward'. No transcripts were detected in mature fruit of either 'Hayward' or *A. chinensis* 'Hort16A'. However a virtual northern analysis based on discovery of related transcripts in libraries in the EST database indicated expression of transcripts in 'Hayward' petals, *A. chinensis* 'Hort16A' vegetative meristems and young fruit of *Vaccinium corymbosum* (blueberry).

**EXAMPLE 7****Expression of multifunctional germacrene D synthase in *Arabidopsis***

- 5 **Preparation of competent *Agrobacterium tumefaciens* GV3101:** *A. tumefaciens*, strain GV3101, containing binary vectors for plant transformation, was inoculated into 10mL of LB or 2YT media containing rifampicin (10 mg/mL); gentamycin (25 mg/mL); and spectinomycin (100 mg/mL) for 24 hours at 28 °C, with shaking at 200 rpm. These cultures were then used to inoculate a further 100-200mL of LB or 2YT media, with
- 10 antibiotics as above, which were again grown for 24 hours at 28 °C, with shaking. The cells were collected by centrifugation (3,500 x g, 10 min, 4 °C) and resuspended, to a final OD<sub>600</sub> of 0.8, in 5% sucrose solution. Silwet L-77 was added to a concentration of 0.05%.
- 15 **Transformation of *Agrobacterium tumefaciens* GV3101:** 45 mL aliquots of competent *Agrobacterium* cells were thawed gently on ice. 50-200 ng of plasmid DNA was added to each aliquot and gently mixed, then 40 mL of the cell/plasmid mixture was pipetted into a pre-chilled electroporation cuvette (0.2 cm gap, Bio-Rad). The cells were electroporated using a BioRad GenePulser, on the following settings:
- 20 Voltage: 2.5 kV  
Capacitance: 25 mFd  
Resistance: 400 Ohms  
The time constant for the pulse was typically 7-9 ms.
- 25 The cells were immediately recovered by addition of 1 mL LB media, then decanted into sterile 15 mL centrifuge tubes and incubated at room temperature, with shaking (60 rpm). After 2 hours, 10 mL and 100 mL of the transformed bacteria was spread onto separate LB plates containing rifampicin (10 mg/mL); gentamycin (25 mg/mL); and spectinomycin (100 mg/mL); then grown for 48 hours at 28-30 °C.

30

**Transformation of *Arabidopsis thaliana* by floral dip:** *Agrobacterium tumefaciens*, containing the appropriate plasmids, were grown in 5mL cultures of LB media

containing rifampicin (10 mg/mL); gentamycin (25 mg/mL); and spectinomycin (100 mg/mL) for 24 hours at 28 °C. The cells were collected by centrifugation and resuspended, to a final OD<sub>600</sub> of 0.8, in 5% sucrose solution. Silwet L-77 was added to a concentration of 0.05%.

5

Healthy *Arabidopsis* plants, around five weeks old, showing a number of immature flower clusters, were used for the dipping procedure. The whole of the aboveground portion of the plant was dipped into the *Agrobacterium* suspension, and gently agitated for 3-5 seconds. The dipped plants were then placed in humidity chambers in reduced light for 2-3 days, before being allowed to flower and set seed as normal. The seed was harvested upon complete drying of the plants (5-6 weeks after dipping).

10

**Growth of seed from transformed *Arabidopsis*:** Approximately 1000 T1 seeds were measured into a microcentrifuge tube. The seed was sterilised by a 1.5% bleach solution containing 0.01% Triton-X, mixed, and incubated for 15 min with occasional mixing. The seed was washed several times with distilled water, and resuspended in 0.1% agarose, prior to plating on 0.5X MS media, containing 100 mg/mL kanamycin. The plates were placed into growth rooms with a 12-hour light/12-hour dark cycle. After 2-3 weeks growth the plants were transferred onto fresh plates to confirm kanamycin resistance and were allowed to continue growing in glasshouse conditions until required for headspace analysis.

15

20

#### Headspace analysis of *Arabidopsis*

*Arabidopsis* plants were grown in soil in a glass house. When mature (8 weeks old) and with inflorescences, three plants were transferred to a 3 L glass vessel with ground glass joint flanges. The vessel was sealed with an appropriate ground glass flanged lid which was fitted with a headspace adaptor with an air inlet and outlet. Volatiles were collected onto 100mg Chromosorb 105 adsorbent traps, which were fitted to the air outlet port of each adaptor. The volatiles were purged from the vessel onto the traps with clean air at 50 mL min<sup>-1</sup> for 6 days. The traps were analysed by GC-MS analysis immediately after extraction of volatiles in the same manner as for the dynamic headspace of kiwifruit flowers.

25

30

## RESULTS

Germacrene D synthase transcripts were detected in transgenic *Arabidopsis* (see Figure 19). There were a number of compounds detected in the headspace of the transformed *Arabidopsis* grown in the growth vessel. In order to clarify if any of these compounds were germacrene D (or other sesquiterpenes) the ions of the base peak (161) and the molecular ion (204) were isolated from the TIC (see Figure 20). This work showed that a number of sesquiterpenes were produced in the *Arabidopsis* including *delta*-elemene, alloaromadendrene, *beta*-caryophyllene, thujopsene, an unidentified sesquiterpene which may be Germacrene D or *gamma*-muurolene, *delta*-cadinene, *gamma*-cadinene, possibly *alpha*-muurolene and elemol. Recently, literature has been published which cites caryophyllene, thujopsene, *beta*-chamigrene, humulene, farnesene, bisabolene, and nerolidol as being present in flowering *Arabidopsis* plants (Aharoni *et al.*, The Plant Cell, Vol. 15, 2866–2884 (2003)). Neither this work or the published literature suggests that *delta*-elemene, Germacrene D, *gamma*-muurolene, *delta*-cadinene, *gamma*-cadinene, *alpha*-muurolene or elemol can be found in wild type flowering *Arabidopsis*. The presence of *delta*-elemene, Germacrene D, *gamma*-muurolene *delta*-cadinene, *gamma*-cadinene, *alpha*-muurolene or elemol in the transformed *Arabidopsis* is consistent with the results shown by the purified recombinant protein.

The above Examples are an illustration of practice of the invention. It will be appreciated by those skilled in the art that the invention can be carried out with numerous modifications and variations. For example, variations to the nucleotide sequences may be used and the sequences may be expressed in different organisms.



## CLAIMS

1. An isolated polynucleotide encoding a multifunctional germacrene-D synthase.
- 5 2. An isolated polynucleotide having the sequence of SEQ ID NO:1 or a fragment or variant thereof encoding a polypeptide with multifunctional germacrene-D synthase activity.
- 10 3. A polynucleotide as claimed in claim 1 or claim 2 wherein the polynucleotide is capable of facilitating the conversion of FDP to a mixture of germacrene-D and one or more other sesquiterpenes selected from *delta*-cadinene, *delta*-elemene, elemol, *gamma*-muurolene, *gamma*-cadinene, *gamma*-elemene and germacrene B.
- 15 4. An isolated polynucleotide as claimed in claim 3 wherein the sequence has at least 60% identity to the nucleotide sequence of SEQ ID NO:1.
5. An isolated polynucleotide as claimed in claim 3 wherein the sequence has at least 90% identity to the nucleotide sequence of SEQ ID NO:1.
- 20 6. An isolated polynucleotide as claimed in claim 3 wherein the sequence has at least 95% identity to the nucleotide sequence of SEQ ID NO:1.
- 25 7. An isolated polynucleotide as claimed in claim 3 wherein the nucleotide sequence is that of SEQ ID NO:1.
8. An isolated polynucleotide encoding the polypeptide of SEQ ID NO:2 or encoding a variant or a fragment of that sequence which has a multifunctional germacrene-D synthase activity.
- 30 9. An isolated polynucleotide as claimed in claim 8 wherein the polypeptide has at least 60% identity with the amino acid sequence of SEQ ID NO:2.

10. An isolated polynucleotide as claimed in claim 8 wherein the polypeptide has at least 90% identity with the amino acid sequence of SEQ ID NO:2.
11. An isolated polynucleotide as claimed in claim 8 wherein the polypeptide has at least 95% identity with the amino acid sequence of SEQ ID NO:2.
12. An isolated polynucleotide as claimed in claim 8 wherein the polypeptide has the sequence of SEQ ID NO:2.
13. An isolated multifunctional germacrene-D synthase polypeptide.
14. An isolated multifunctional germacrene-D synthase having the sequence of SEQ ID NO:2 or a fragment or variant thereof with multifunctional germacrene-D synthase activity.
15. An isolated multifunctional germacrene-D synthase as claimed in claim 14 wherein the amino acid sequence has at least 60% identity with the sequence of SEQ ID NO:2.
16. An isolated multifunctional germacrene-D synthase as claimed in claim 14 wherein the amino acid sequence has at least 90% identity with the sequence of SEQ ID NO:2.
17. An isolated multifunctional germacrene-D synthase as claimed in claim 14 wherein the amino acid sequence has at least 95% identity with the sequence of SEQ ID NO:2.
18. An isolated multifunctional germacrene-D synthase as claimed in claim 14 wherein the amino acid sequence is a mature sequence derived from SEQ ID NO:2.
19. A genetic construct comprising a polynucleotide of any one of claims 1 to 12.

20. A genetic construct comprising an open reading frame polynucleotide encoding a polypeptide of any one of claims 13 to 18.

21. A genetic construct as claimed in claim 20 further comprising a promoter  
5 sequence.

22. A genetic construct as claimed in claim 21 which further comprises a termination sequence.

10 23. A genetic construct as claimed in claim 22 wherein the sequence of the encoded polypeptide has the amino acid sequence of SEQ ID NO:2 or a fragment thereof with multifunctional germacrene-D activity.

24. A genetic construct comprising in the 5'-3' direction a polynucleotide which  
15 hybridizes to a polynucleotide encoding a polypeptide of any one of claims 13 to 18.

25. A genetic construct as claimed in claim 24 further comprising a promoter sequence.

20 26. A genetic construct as claimed in claim 25 which comprises a termination sequence

27. A genetic construct as claimed in claim 26 wherein the sequence of the encoded polypeptide has the amino acid sequence of SEQ ID NO:2 or a fragment thereof with  
25 multifunctional germacrene-D activity.

28. A vector comprising a genetic construct of any one of claims 19 to 27.

29. A host cell comprising a genetic construct of any one of claims 19 to 27.  
30

30. A transgenic plant cell which includes a genetic construct of any one of claims 19 to 27.

31. A transgenic plant comprising a plant cell as claimed in claim 30.

32. A method of preparing germacrene-D, *delta*-cadinene, *gamma*-cadinene,  
5 *gamma*-muurolene, *gamma*-elemene, *delta*-elemene, elemol or germacrene B  
comprising the steps of

(a) culturing a cell which has been genetically modified with a polynucleotide any  
one of claims 1-12 to provide increased multifunctional germacrene-D synthase activity;

10 (b) providing the cell with farnesyl diphosphate or geranyl diphosphate if  
necessary; and

(c) separating the germacrene-D and/or *delta*-cadinene and/or *delta* elemene and/or  
elemol and/or germacrene B, and/or *gamma*-cadinene, and/or *gamma*-muurolene, and/or  
*gamma*-elemene produced.

15 33. A method for modulating the Germacrene-D and/or *delta*-cadinene and/or  
germacrene B and or elemol and/or *delta*-elemene, and/or *gamma*-cadinene, and/or  
*gamma*-muurolene, and/or *gamma*-elemene production of a plant, the method  
comprising: increasing or decreasing expression of multifunctional germacrene-D  
synthase wherein said increasing or decreasing is achieved by genetic modification to  
20 alter the expression of a gene encoding a multifunctional germacrene-D synthase.

34. A method as claimed in claim 33 wherein the synthase comprises a synthase  
with the sequence of SEQ ID NO: 2.

25 35. A method for modulating germacrene-D and/or *delta*-cadinene and/or  
germacrene B and or elemol and/or *delta*-elemene, and/or *gamma*-cadinene, and/or  
*gamma*-muurolene, and/or *gamma*-elemene production in a plant, the method  
comprising of:

(a) introducing into the plant, a genetic construct of claims 19-27; and

30 (b) transcriptionally expressing the polynucleotide in the plant.

36. A method for modulating germacrene-D and/or *delta*-cadinene and/or germacrene B and or elemol and/or *delta*-elemene, and/or *gamma*-cadinene, and/or *gamma*-muurolene, and/or *gamma*-elemene production in a plant, the method comprising of

- 5 (a) introducing into the plant, a DNA genetic construct of claims 19-27; and  
(b) expressing the polypeptide in the plant.

37. A polynucleotide having at least 15 contiguous nucleotides from SEQ ID NO: 1

10 38. A method of selecting a plant with altered germacrene-D and/or *delta*-cadinene and/or germacrene B and or elemol and/or *delta*-elemene, and/or *gamma*-cadinene, and/or *gamma*-muurolene, and/or *gamma*-elemene content comprising the steps of:

- (a) contacting polynucleotides from at least one plant with at least one polynucleotide comprising at least 15 contiguous nucleotides of the polynucleotide of claim 1 to assess the expression of multifunctional germacrene-D synthase; and  
15 (b) selecting a plant showing altered expression.

39. A method as claimed in claim 38 wherein the polynucleotide has at least 15 contiguous nucleotides from a sequence selected from SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 4 and SEQ ID NO: 7 and the plant is a plant of the genus *Actinidia*.  
20

40. A method as claimed in claim 38 wherein the plant is a plant of the genus *Vaccinium*.

25 41. A method for preparing a sesquiterpene comprising:

- (a) obtaining a polypeptide as claimed in any one of claims 13-18; and  
(b) incubating farnesyl diphosphate in the presence of the polypeptide, and  
(c) separating the germacrene-D and/or *delta*-cadinene and/or germacrene B and or elemol and/or *delta*-elemene, and/or *gamma*-cadinene, and/or *gamma*-muurolene, and/or *gamma*-elemene produced.  
30


42. A method as claimed in claim 41 wherein the products of step (b) are trapped in a matrix providing an acid environment.

43. A method as claimed in claim 42 wherein the matrix is a silica base matrix.



5

44. A method as claimed in claim 41 wherein the sesquiterpene is germacrene D.




1/22





Germacrene D

 $\delta$ -elemene


Elemol

 $\delta$ -cadinene $\gamma$ -cadinene

Germacrene B


 $\beta$ -elemene $\gamma$ -elemene**FIGURE 1**

2/22

**FIGURE 2A**



3/22

**FIGURE 2B**

4/23

1   GTGAAAACCTA AATATGCCCC AGTGTGTATGG TTCAATCTCTA GTTTTCTCTT TTAATTTAAT TCTTCAACCC AGAAAAAATA CATGCAACCTA CCTGTGTGCTC  
 101   AAGCTTTTCCC AATACCAAACT GTTACAAACCA CCACATAGAT TGAACACCCA CATGTAAACGC GTGCGCTCTGC AATTTATCAT CCTATGCAATTT GGGGAGATCA  
 201   TTTCCCTGCC TACTCTTCCG ATGCTATGGG AGAAGAGGTT ATTAACATGG AACAAACAACA ACGACTTCAT CACTGTAAAC AATAGGTGAG AAAAAATGCTA  
 301   GAGGCAAGCTG CTGAAACAATC TTCAACAGATG CTGAACCTTG TCGACAAAAA CCAACGCTTA GCGGTGTCTT ACCATTTTGA AACGTAGATC GAATCAAGCTT  
 401   TACGGCACAT ATACAAAACC TGTGATATAC AATTTGATGA TCTCCACACT GCTGCTCTCT CTTTGCGTT ACTTGAACAA CAAGGATATC CAGTTTCTTG  
 501   TGAATATGTT GACAAAATCA AAGAACGCAA AGGTGAGTTT CAAGAAATCCA TAATCAAGCA TGTGCAAGGA ATGTAAAGTT TGTATGAAAG TACATGTCTA  
 601   AGGATACACG GAGAAAGATAT ACTAGACGAA GCACATAGCT TTACCATCAC TCAACTTCGG TCCGCATTTGC CCAACTTAAG CACTGCTTTT AAGGAAACAA  
 701   TCATTTCAAT TCTGAAACCA CCCCATCCACA AGGGGTGAG AACGCTCAAC GCAAGGAGCC ACAATTTAT TTTTGAACAG AATGATTTCC CACTGCAAAAG  
 801   CCTTTTGAAT TTGCGAAATAT TAGATTTCAA CTTATTTCAA AAGTTGCACC AGAGGAGGCT ATGTGAAATC ACAAGGTTGG GGAAGAATTT GAAATTTTCCA  
 901   AAGACACTAC CTTTTCGCA GAGACGAAATG GTAGAGTGTCT ACTTTTGAAT ACTTGGGTTG TACTTTGAGC CCGAATATCT GCTTGTATGG AGGATGCTAA  
 1001   CCAAAGGTGAT TGGCATGATTT TCCATTTATCG ATGACATCTA CGATGTCTAC GGTACTTTGG AAGAACTTGT TCTCTTCACT GATGCAATTT AGAGGTGGGA  
 1101   GATCAATGCC TTGATCAAC TTCCAGAGTA TAGGAAACCTA TGTATATCA GACTTTTGGG TGTATATAGT ATGATTTGATG AAGAGATTTGC GAAAGCAAGGA  
 1201   AGATCTTAT TCGGTAGACTA TGCAAAATCT TCAATGAAA TTTTGGTTAG AGCATATCTC GAAAGAAAGCA AATGTTTTCA CCAAGGATAT GTTCCAACTA  
 1301   TGGAAAGATA TAGGCAAGTT GCATTTAGTAA CCGCGGGTTA CAAAATGCTT CAAATCTCTT CCTTTGTTGG CATGGAGAT TTGCGCAACCA AAGAGGCTTT  
 1401   TGAATGGGTT TCAAAATGATC CTTTAAATTT TCAAGCTGCA TCAATGATAG CCAAGCTCAA GATGTGCAAT GTTGGCCACA AGTTTGAACA AATGAGAGGG  
 1501   CACGTGGCGT CGGCTGTGGA ATGCTACAGT AAGCAACATG GTACAAACGA GGAAGAGGCT ATTAATTTGAT TGGATTAACA AGTTACACAT TCAATGGAAAG  
 1601   ACATCAACGC AGAGTGCCCTC TGGCCAAATCA AGGTCCCAAT GCTCTCTCTT CCGCGAGTTT TCAATCTTGC ACGAGTGTCT TATGTTATAT ACCAGGATGA  
 1701   AGACGGGATC ACTCATCTTG GAAACCAAGGT CGAGAACTTT GTAACTCTAG TGTATATCGA TTCTATGCCA ATCAATTAGA AATGTAAACA AGACACTGAA  
 1801   GTGGAGGCAAT AAAATAAATC AATAGTTGAT TTAAGGTGAG GTTAGGTGAG GGGGATTTCTT ACCATTAAGA GATATTTCTTG CTAATAAGCA ATTAATTCAA  
 1901   TGCATTTTCCA ATAAATTAAT TTAGCCAGTT GTTCTCTATC TGTGTTTTT TTGTGTTCTT TTTCCTTCTT AATATATAAA TTAATTTAAT TTGGCAAAAA  
 2001   AAAAAATAA AAAAAATA

FIGURE 3

5/22

1 MQLPCAQALP IPTVTTTTSI EPPHVTRRSA NYHPSIWGDH FLAYSSDAME  
51 EEVINMEQQQ RLHHLKQKVR KMLEAAAEQS SQMLNLVDKI QRLGVSYHFE  
101 TEIETALRHI YKTCDYHFDD LHTAALSFRL LRQQGYPVSC DMFDKFKNSK  
151 GEFQESIISD VQGMLSLYEA TCLRIHGEDI LDEALAFITIT QLRSAIPNLS  
201 TPFKEQIIHA LNQPIHKGLT RLNARSHILF FEQNDCHSKD LLNFAKLDEF  
251 LLQKLHQREL CEITRWKDL NFAKTLPEAR DRMVECYFWI LGVYFEPQYL  
301 LARRMLTKVI AMISIIDDIY DVYGTLEELV LFTDAIERWE ISALDQLPEY  
351 MKLCYQALLD VYSMIDEEMA KQGRSYCVDY AKSSMKILVR AYFEEAKWFH  
401 QGYVPTMEEY MQVALVTAGY KMLATSSFVG MGDLATKEAF DWVSNDPLIV  
451 QAASVIGRLK DDIVGHKFEQ KRGHVASAVE CYSKQHGTTE EEAIIELDKQ  
501 VTHSWKDINA ECLCPIKVPM PLLARVLNLA RVLYVIYQDE DGYTHPGTKV  
551 ENFVTSVLID SMPIN\*

**FIGURE 4**

**6/22**

[illegible]

## FIGURE 5

7/22

>EST 80968 (*A. deliciosa* variant of multifunctional germacrene D synthase)

CTAAATAGGCCAAGTGTGTAGGTTTCATCTCTAGTTTTTCTCTTGAAAACATAAATAGGCCAAGTG  
TGTAGGTTTCATCTCTAGTTTTTCTCTTTAAATTAATCCTTCAACCCAGAAAAAACATGCAACTA  
CCTTGTGCTCAAGCTTTGCCAATACCAACTGTTACAACCAACTAGTATTGAACCACCACATGTA  
ACTCGTGCATCTGCAAATTATCATCCTAGCATTGGGGAGATCATTTCTCGCCTACTCTTCCGAT  
GCTATGGAAGAAGAGGATATTAACATGGAACAACAACGACTTCATCACCTGAAACAAAAGGTG  
AGAAAAATGCTAGAGGCAGCTGCTAAACAATCTTCACAGATGCTGAACCTCGTCGACAAAATCCAA  
CGCTTAGGCGTGTCTTACCATTTTGAAGTGAAGATCGAAACAGCTTTACGGCACATATACAAAACC  
TGTGATTACCATTTTGATGATCTCCACACTGCTGCTCTCTCTTTTCGGTTACTTAGACAACAAGGA  
TATCCAGTTTCTTGTGACATGTTCCGCAAATTCAAGAACTGCAAAGGTGAGTTTCAAGAATCCATA  
ATCAGCGATGTGCGAGGAATGTTAAGCTTGATGAAGCTACATGTCTAAGGATACGCGGAGAAGAT  
ATACTAGACGAAGCACTAGCTTTTACCACGACTCAGCTTCAGTCTGCATTGCCCAACTTAAGCACT  
CCTATCAAGGAACAAATCATTCATGCTCTGAACCCAGCCCATCCACAAGTGGTTGACAAGGCTCGAC  
GCAAGGCGCCACATTTTATTCTTGAACAGAATGATTGCCATGGCAAAGACCTTTTGAATTTTCGCA  
AAATTAGATTTCAACTCGTTACAAAAGTTGCACCAGAGGGAGCTATGTGAAATCACAAGGTGGTGG  
AAAGATCTGGATTTTGCCAAGAACTACCTTTTGCCAGAGACAGAATGGTAGAGTGCTACTTCTGG  
ATACTTGGGGTGTACTTTGAGCCCCAATATTTGCGTGCTAGGAGGATGCTAACCAAGGTGATTGCC  
TTGACTTCCATTATCGATGACATCTACGATGTCTACGGTACCTTGAAGAAGCTTGTCTCTTCACT  
GATGCAATTGAGAGGTGGGAAATTAGTGCCTTGGATAACCTTCCAGATTATATGAAACTATGTTAT  
CAAGCACTTTTGATGTTTATAGTATGATTGATGAAGAGATGGCCAAGCAAGGAAGATCTTATTGC  
GTAGACTATGCAAAATCTTCAATGAAAATTTTGGTTAGAGCATACTTCAAGAAGCCAAATGGTTT  
CACCAAGGATATGTTCCAACTATGGAAGAGTATATGCAAGTTGCATTAGTAACCGCGGGTTACAAA  
ATGCTTGCAACCTCTTCTTTGTTGGCATGGGAGAGTTGGCAACCAAGAGGCCCTTTGATTGGGTG  
TCAAATGATCCTTTAATTGTTCAAGCTGCATCAGTGATAGGCAGACTCAAGGATGACATTTGTTGGC  
CACAAGTTTGAGCAAAAGAGAGGGCACGTGGCGTCGGCTGTGCAATGCTACAGTAAGCAACATGGT  
ACAATAGAGGAAGAGGCTATTATTGAATTGGATAAACAAGTTACACATTATGGAAGACATCAAC  
GCAGAGTGCTCTGCCAATCAAGGTCCCAATGCCTCTTCTTGCGCGAGTTCTCAATCTTGACGGA  
GTGCTTTATGTTATATACAGGATGAAGACGGCTACACTCATTCTGGAACCAAGGTCAAGAACTTT  
GCAACCTCAGTGCTTATCGATTCTATGCCAATCAATTAGAAAATGTAACAAGACTGAAGTGGAG  
GCATAAATAAATTCAAAGTTGGCTTAAAGTTGGGCTAAAAA

>EST 304951 (*A. chinensis* variant of multifunctional germacrene D synthase)

ATCTTATTGCGTAGACTATGCAAAATCTTCAATGAAAAGTTTGGTTAGAGCATACTTCAAGAAGC  
CAAATGGTTTCACCAAGGATATGTTCCAATGGAAGAGTATATGCAAGTTGCAATAGTAACCGG  
GGCTTACAAAATCTTGAACCACTTCTTTGTTGGCATGGGAGAGTTGGCAACCAAGAGGTCTT  
TGATTGGGTGTCAAATGATCCTTTAATTGTTCAAGCTGCATCAATTGTTCCAGACTCACGGATGA  
CATTGTTGGCCACAAGTTTGAGCAAAAGAGAGGGCACGTGGCATCGGCGGTTGAATGCTACATGAA  
GCAACATGGTACAACAGAGGAAGAGGCCATTGTTGAATTGTATAAGCAAGTTACAAATGCTAGGAA  
AGACATGAATGCAGAGTGCTCTTCCCCACCAAGGTCCCAATGCCTCTTCTCGTGAGAGTTCTCAA  
TCTTGACAGAGTGATTAATGTTCTATACAAGGATGAAGATGGCTACACTCATTCAAGAACCAAGGT  
TAAGAAATTTGTGACCTCAGTGCTTGTAGATTTTGTGCCGATCAGCTAGCAAACGTTCTCTCTAC  
CACATGTTAATTAGTCTGCTTGTAAATGCAGTTTACTAATATGAAATTTAATAAATGCGTATTTTC  
CAATAAAGGAATTTAAAAA

>EST 82293 (*Vaccinium corymbosum* variant of multifunctional germacrene D synthase)

GGAAGCCAAATGGTTTCATGAAGGTTATGTTCCGAGTATGGAAGAGTATATGAGAGTTGCACTGGT  
TACCGGTGCTTACAAAATGCTTGCAACCACTTCTTTTGTGGCATGGGGGATTGTTGACCAAGA  
GGCCTTTGAATGGGTGTCAAGTGATCCTTTAATTGTTGAAGCTGCATCCGTGATTGTCAGACTCAT  
GGATGATATGGCAGGCCACAAGTTTGAGCAAGAGAGAGGACACGTGGCTTCGGCAGTTGAATGCTA  
CATGAAACAACATGGTGCAACACAAGAAGTGGTTCTTCTTGAATTTAAAAAAGAGTTACAAATGC  
ATGGAAGACATGAACGCAGAGTGCTTCCGCCCACTGCCGTTCATGCCTCTCTCACCAGGAT  
TCTCAATCTCGCACGAGTGATCAATGTTATATACAAGGATGAAGATGGGTACACTCATTCTGGAAC  
AAAGCTCAAGAAGCTTTGTAATCTCAGTGCTTATCAGATTCTGTGCGGATCAATTAGCAAACAGTAGT  
CCTAACTTAAATAATCTGTTGGCTTATAACTTTATAAGTGTGCTGAAATGTTCTAGTGAAGTGGT  
AAGGATGTATTTCCGATATGTAGCTCTATCTCCACTGTACGGTTGTAATCTTGCTCTCTTCTACTA  
AGAAAGCTCATTAATCGCTGCTTAAATGTAAAGCCAAGTGTCTCAAGTTTATCGTCAAACAAGTT  
CTGTTTACGATTTTGTGGA

**FIGURE 5 (continued)**

SUBSTITUTE SHEET (RULE 26)

8/22

&gt;72838

MQLPCAQALPIPTVTTNTSIEPPHVTRRSANYHPSIWGDHFLAYSSDAMEEEDINMEQQQRLHHLK  
 QKVRKMLEAAAEQSSQMLNLVDKIQRLGVSYHFETEIETALRHIYKTCDYHFDDLHTAALSFRLLR  
 QQGYPVSCDMFDKFKNSKGEFQESIISDVRGMLSLYEATCLMIHGEDIIDEALAFITITQLRSALPN  
 LSTPFKEQIIHALNQPIHKGLTRLNARSHILFFEQNDCHSKDLLNFAKLDFNLLQKLHQRELCEIT  
 R~~~~~  
 ~~~~~~EISALDQLPEYMKLCYQALLDVYSMIDEEMAKQGRSYCVDYAKSSMKILVRAYFEEA  
 KWFHQGYVPTMEEYMQVALVTAGYKMLATSSFVGMGELATKEAFDWVSNPLIVQAASVIGRLKDD  
 IVGHKFEQKRGHVASAVECYSKQHGTTEEEAIIELYKQVTHSWKDMNAECLCPTKVPMPLLARVLN  
 LARVLYVIYQDADGYTHSGTKVKNFVTSVLIDSMPIN

&gt;80968

MQLPCAQALPIPTVTTNTSIEPPHVTRRSANYHPSIWGDHFLAYSSDAMEEEDINMEQQQRLHHLK  
 QKVRKMLEAAAKQSSQMLNLVDKIQRLGVSYHFETEIETALRHIYKTCDYHFDDLHTAALSFRLLR  
 QQGYPVSCDMFQKFKNCKGEFQESIISDVRGMLSLYEATCLRIRGEDILDEALAFITITQLQSALPN  
 LSTPIKEQIIHALNQPIHKWLTRLDARRHILFFEQNDCHGKDLLNFAKLDFNSLQKLHQRELCEIT  
 RWWKDLDFAKKLPFARDRMVECYFWILGVYFEPQYLRARRMLTKVIALTSIIDDIYDVYGTLEELV  
 LFTDAIERWEISALDNLPDYMKLCYQALLDVYSMIDEEMAKQGRSYCVDYAKSSMKILVRAYFEEA  
 KWFHQGYVPTMEEYMQVALVTAGYKMLATSSFVGMGELATKEAFDWVSNPLIVQAASVIGRLKDD  
 IVGHKFEQKRGHVASAVECYSKQHGTIEEEAIIELDKQVTHSWKDMNAECLCPIKVPMPLLARVLN  
 LARVLYVIYQDEDGYTHSGTKVKNFATSVLIDSMPIN

&gt;304951

YCVDYAKSSMKSLVRAYFEEAKWFHQGYVPTMEEYMQVAIVTGAYKILATTSFVGMGELATKEVFD  
 WVSNDPLIVQAASIVSRLTDDIVGHKFEQKRGHVASAVECYMKQHGTTEEEAIIELYKQVTNAWKD  
 MNAECLFPTKVPMPLLVRLNLARVINVLYKDEDGYTHSRTKVKKFVTSVLVDFVPIS

&gt;82293

EAKWFHEGYVPSMEEYMRVALVTGAYKMLATTSFVGMGDLVTKEAFEWVSSDPLIVEAASVICRLM  
 DDMAGHKFEQERGHVASAVECYMKQHGTQEVVLEFKKRVTNNAWKDMNAECLRPTAVPMPLLTRV  
 LNLARVINVIYKDEDGYTHSGTKLKNFVISVLIDSVPIN

## FIGURE 6

9/22




FIGURE 7

10/22




FIGURE 8



11/22




FIGURE 9

12/22




FIGURE 10

13/22




FIGURE 11

14/22




FIGURE 12

15/22




FIGURE 13

16/22




FIGURE 14

17/22

# EST 75565 Temperature Response




FIGURE 15

18/22

# EST 75565 Metal Ion Responses




FIGURE 16



19/22






FIGURE 17

20/22

**FIGURE 18**

21/22

**FIGURE 19**

22/22




FIGURE 20

497047kxr.ST25.txt  
SEQUENCE LISTING

<110> THE HORTICULTURE AND FOOD RESEARCH INSTITUTE OF NEW ZEALAND LIMITED

<120> ENZYMES AND POLYNUCLEOTIDES ENCODING THE SAME

<130> 497047 KXR

<150> NZ 523384

<151> 2002-12-24

<160> 16

<170> PatentIn version 3.1

<210> 1

<211> 2019

<212> DNA

<213> Actinidia deliciosa

<400> 1  
gtgaaaacta aaataggcca agtgtgtagg ttcattctcta gtttttctct ttaaattaat 60  
tcttcaaccc agaaaaaaaa catgcaacta ccttgtgtgc aagctttgcc aataccaact 120  
gttacaacca ccaactagtat tgaaccacca catgtaacgc gtcgctctgc aaattatcat 180  
cctagcattt ggggagatca tttcctcgcc tactcttccg atgctatgga agaagagggtt 240  
attaacatgg aacaacaaca acgacttcat cacctgaaac aaaagggtgag aaaaatgcta 300  
gaggcagctg ctgaacaatc ttcacagatg ctgaacctcg tcgacaaaat ccaacgctta 360  
ggcgtgtctt accattttga aactgagatc gaaacagctt tacggcacat atacaaaacc 420  
tgtgattacc attttgatga tctccacact gctgtctctc cttttcggtt acttagacaa 480  
caaggatatc cagtttcttg tgatatgttc gacaaattca agaacagcaa aggtgagttt 540  
caagaatcca taatcagcga tgtgcaagga atgttaagtt tgtatgaagc tacatgtcta 600  
aggatacacg gagaagatat actagacgaa gcactagctt ttaccatcac tcaacttcgg 660  
tccgcattgc ccaacttaag cactcctttc aaggaacaaa tcattcatgc tctgaaccag 720

497047kxr.ST25.txt

```

cccatccaca aggggttgac aaggctcaac gcaaggagcc acatTTtatt ttttgaacag      780
aatgattgcc atagcaaaga ctttttgaat ttcgcaaaat tagatttcaa cttattacaa      840
aagttgcacc agaggagct atgtgaaatc acaagggtgt ggaaagattt gaattttgca      900
aagacactac cttttgccag agacagaatg gtagagtgt acttttggat acttgggggtg      960
tactttgagc cccaatatct gcttgctagg aggatgctaa ccaagggtgat tgccatgatt     1020
tccattatcg atgacatcta cgatgtctac ggtaccttgg aagaacttgt tctcttcact     1080
gatgcaattg agagggtgga gatcagtgcc ttggatcaac ttccagagta tatgaaacta     1140
tgttatcaag cacttttga tgtttatagt atgattgatg aagagatggc gaagcaagga     1200
agatcttatt gcgtagacta tgcaaaatct tcaatgaaaa ttttgggttag agcatacttc     1260
gaagaagcca aatggtttca ccaaggatat gttccaacta tggaagagta tatgcaagtt     1320
gcattagtaa ccgcgggtta caaatgctt gcaacctctt cctttgttgg catgggagat     1380
ttggcaacca aagaggcctt tgattgggtg tcaaatgatc ctttaattgt tcaagctgca     1440
tcagtgatag gcagactcaa ggatgacatt gttggccaca agtttgagca aaagagaggg     1500
cacgtggcgt cggctgtcga atgctacagt aagcaacatg gtacaacaga ggaagaggct     1560
attattgaat tggataaaca agttacacat tcatggaaag acatcaacgc agagtgcctc     1620
tgcccaatca aggtcccaat gcctcttctt gcgcgagttc tcaatcttgc acgagtgtt     1680
tatgttatat accaggatga agacggatac actcatcctg gaaccaaggt cgagaacttt     1740
gtaacctcag tgcttatcga ttctatgcca atcaattaga aaatgtaaca agacactgaa     1800
gtggaggcat aaataaattc aaaagttgat ttaaagttgg ggtagtgaac ggggattctt     1860
accattaaga gatattcttg ctaaaaagca attaatcaa tgcatttcca ataaaataat     1920
ttagccagtt gttcttcac tttgtttttt tttgtttctc tttcctttct aaatataaaa     1980
ttataattaa ttggcaaaaa aaaaaaaaaa aaaaaaaaaa                        2019

```

&lt;210&gt; 2

&lt;211&gt; 565

&lt;212&gt; PRT

&lt;213&gt; Actinidia deliciosa

&lt;400&gt; 2

```

Met Gln Leu Pro Cys Ala Gln Ala Leu Pro Ile Pro Thr Val Thr Thr
1          5          10          15

```

```

Thr Thr Ser Ile Glu Pro Pro His Val Thr Arg Arg Ser Ala Asn Tyr
          20          25          30

```

497047kxr.ST25.txt

His Pro Ser Ile Trp Gly Asp His Phe Leu Ala Tyr Ser Ser Asp Ala  
 35 40 45  
 Met Glu Glu Glu Val Ile Asn Met Glu Gln Gln Gln Arg Leu His His  
 50 55 60  
 Leu Lys Gln Lys Val Arg Lys Met Leu Glu Ala Ala Ala Glu Gln Ser  
 65 70 75 80  
 Ser Gln Met Leu Asn Leu Val Asp Lys Ile Gln Arg Leu Gly Val Ser  
 85 90 95  
 Tyr His Phe Glu Thr Glu Ile Glu Thr Ala Leu Arg His Ile Tyr Lys  
 100 105 110  
 Thr Cys Asp Tyr His Phe Asp Asp Leu His Thr Ala Ala Leu Ser Phe  
 115 120 125  
 Arg Leu Leu Arg Gln Gln Gly Tyr Pro Val Ser Cys Asp Met Phe Asp  
 130 135 140  
 Lys Phe Lys Asn Ser Lys Gly Glu Phe Gln Glu Ser Ile Ile Ser Asp  
 145 150 155 160  
 Val Gln Gly Met Leu Ser Leu Tyr Glu Ala Thr Cys Leu Arg Ile His  
 165 170 175  
 Gly Glu Asp Ile Leu Asp Glu Ala Leu Ala Phe Thr Ile Thr Gln Leu  
 180 185 190  
 Arg Ser Ala Leu Pro Asn Leu Ser Thr Pro Phe Lys Glu Gln Ile Ile  
 195 200 205  
 His Ala Leu Asn Gln Pro Ile His Lys Gly Leu Thr Arg Leu Asn Ala  
 210 215 220  
 Arg Ser His Ile Leu Phe Phe Glu Gln Asn Asp Cys His Ser Lys Asp  
 225 230 235 240  
 Leu Leu Asn Phe Ala Lys Leu Asp Phe Asn Leu Leu Gln Lys Leu His  
 245 250 255  
 Gln Arg Glu Leu Cys Glu Ile Thr Arg Trp Trp Lys Asp Leu Asn Phe  
 260 265 270  
 Ala Lys Thr Leu Pro Phe Ala Arg Asp Arg Met Val Glu Cys Tyr Phe  
 275 280 285

497047kxr.ST25.txt

Trp Ile Leu Gly Val Tyr Phe Glu Pro Gln Tyr Leu Leu Ala Arg Arg  
 290 295 300  
 Met Leu Thr Lys Val Ile Ala Met Ile Ser Ile Ile Asp Asp Ile Tyr  
 305 310 315 320  
 Asp Val Tyr Gly Thr Leu Glu Glu Leu Val Leu Phe Thr Asp Ala Ile  
 325 330 335  
 Glu Arg Trp Glu Ile Ser Ala Leu Asp Gln Leu Pro Glu Tyr Met Lys  
 340 345 350  
 Leu Cys Tyr Gln Ala Leu Leu Asp Val Tyr Ser Met Ile Asp Glu Glu  
 355 360 365  
 Met Ala Lys Gln Gly Arg Ser Tyr Cys Val Asp Tyr Ala Lys Ser Ser  
 370 375 380  
 Met Lys Ile Leu Val Arg Ala Tyr Phe Glu Glu Ala Lys Trp Phe His  
 385 390 395 400  
 Gln Gly Tyr Val Pro Thr Met Glu Glu Tyr Met Gln Val Ala Leu Val  
 405 410 415  
 Thr Ala Gly Tyr Lys Met Leu Ala Thr Ser Ser Phe Val Gly Met Gly  
 420 425 430  
 Asp Leu Ala Thr Lys Glu Ala Phe Asp Trp Val Ser Asn Asp Pro Leu  
 435 440 445  
 Ile Val Gln Ala Ala Ser Val Ile Gly Arg Leu Lys Asp Asp Ile Val  
 450 455 460  
 Gly His Lys Phe Glu Gln Lys Arg Gly His Val Ala Ser Ala Val Glu  
 465 470 475 480  
 Cys Tyr Ser Lys Gln His Gly Thr Thr Glu Glu Glu Ala Ile Ile Glu  
 485 490 495  
 Leu Asp Lys Gln Val Thr His Ser Trp Lys Asp Ile Asn Ala Glu Cys  
 500 505 510  
 Leu Cys Pro Ile Lys Val Pro Met Pro Leu Leu Ala Arg Val Leu Asn  
 515 520 525  
 Leu Ala Arg Val Leu Tyr Val Ile Tyr Gln Asp Glu Asp Gly Tyr Thr  
 530 535 540



497047kxr.ST25.txt

His Pro Gly Thr Lys Val Glu Asn Phe Val Thr Ser Val Leu Ile Asp  
 545 550 555 560

Ser Met Pro Ile Asn  
 565

&lt;210&gt; 3

&lt;211&gt; 1815

&lt;212&gt; DNA

&lt;213&gt; Actinidia deliciosa

&lt;400&gt; 3

```

ctaaaatagg ccaagtgtgt aggttcatct ctagtttttc tcttgaaaac taaaataggc      60
caagtgtgta ggttcatctc tagtttttct cttgaaaact aaaataggcc aagtgtgtag      120
gttcatctct agtttttctc tttaaattaa tccttcaacc cagaaaaaaa acatgcaact      180
accttgtgct caagctttgc caataccaac tgttacaacc aacactagta ttgaaccacc      240
acatgtaact cgtcgatctg caaattatca tcctagcatt tggggagatc atttcctcgc      300
ctactcttcc gatgctatgg aagaagagga tattaacatg gaacaacaac aacgacttca      360
tcacctgaaa caaaagggtga gaaaaatgct agaggcagct gctgaacaat cttcacagat      420
gctgaacctc gtcgacaaaa tccaacgctt aggcgtgtct taccattttg aaactgagat      480
cgaaacagct ttacggcaca tatacaaaac ctgtgattac cattttgatg atctccacac      540
tgctgctctc tcttttcggt tacttagaca acaaggatat ccagtttctt gtgatatgtt      600
cgacaaattc aagaacagca aagggtgaatt tcaagaatcc ataatcagcg atgtgcgagg      660
aatgttaagt ttgtatgaag ctacatgtct aatgatacac ggagaagata tactagacga      720
agcactagct ttaccatca ctcaacttcg gtccgcattg cccaacttaa gcactccttt      780
caaggaacaa atcattcatg ctctgaacca gcccatccac aaggggttga caaggctcaa      840
tgcaaggagc cacatthttat tttttgaaca gaatgattgc catagcaaag accttttgaa      900
tttcgcaaaa ttagatttca acttattaca aaagttgcac cagagggagc tatgtgaaat      960
cacaagggtga gatcagtgcc ttggatcaac ttccagagta tatgaaacta tgttatcaag     1020
cacttttgga tgthttatagt atgattgatg aagagatggc gaagcaagga agatcttatt     1080
gcgtagacta tgcaaaatct tcaatgaaaa ttttggttag agcatacttc gaagaagcca     1140
aatggtttca ccaaggatat gttccaacta tggaagagta tatgcaagtt gcattagtaa     1200
ccgcgggtta caaaatgctt gcaacctctt cctttgttgg catgggagag ttggcaacca     1260
aagaggcctt tgattgggtg tcaaatgatc ctttaattgt tcaagctgca tcagtgatag     1320

```

## 497047kxr.ST25.txt

gcagactcaa ggatgacatt gttggccaca agtttgagca aaagagaggg cacgtggcgt 1380  
cggtgttga atgctacagt aagcaacatg gtacaacaga ggaagaggct attattgaat 1440  
tgtataaaca agttacacat tcatggaaag acatgaacgc agagtgcctc tgcccaacca 1500  
aggtcccaat gcctcttctt gcgcgagttc tcaatcttgc acgagtgtt tatgttatat 1560  
accaggatgc agatggctac actcattctg gaaccaaggc caagaacttt gtaacctcag 1620  
tgcttatcga ttctatgcca atcaattaga aaatttaaca agacactgaa gtggagggtat 1680  
aaataaatc aaaagttgat ttaaagttgg gctagtgaac ggggattctt accattaaga 1740  
gatattcttg ctaaaaagca attaattcaa tgcatttcca ataaaataat ttagccagct 1800  
gttgttcaaa aaaaa 1815

&lt;210&gt; 4

&lt;211&gt; 1897

&lt;212&gt; DNA

&lt;213&gt; Actinidia deliciosa

&lt;400&gt; 4

ctaaaatagg ccaagtgtgt aggttcatct ctagtttttc tcttgaaaac taaaataggc 60  
caagtgtgta gggtcatctc tagtttttct ctttaaatta atccttcaac ccagaaaaaa 120  
aacatgcaac taccttgtgc tcaagctttg ccaataccaa ctgttacaac caacactagt 180  
attgaaccac cacatgtaac tcgtcgatct gcaaattatc atcctagcat ttggggagat 240  
catttcctcg cctactcttc cgatgctatg gaagaagagg atattaacat ggaacaacaa 300  
caacgacttc atcacctgaa acaaaagggtg agaaaaatgc tagaggcagc tgctaaacaa 360  
tcttcacaga tgctgaacct cgtcgacaaa atccaacgct taggcgtgtc ttaccatttt 420  
gaaactgaga tcgaaacagc ttacggcac atatacaaaa cctgtgatta ccattttgat 480  
gatctccaca ctgctgctct ctcttttcgg ttacttagac aacaaggata tccagtttct 540  
tgtgacatgt tcggcaaatt caagaactgc aaagggtgagt ttcaagaatc cataatcagc 600  
gatgtgcgag gaatgttaag cttgtatgaa gctacatgtc taaggatacg cggagaagat 660  
atactagacg aagcactagc ttttaccacg actcagcttc agtctgcatt gcccaactta 720  
agcactccta tcaaggaaca aatcattcat gctctgaacc agcccatcca caagtgggtg 780  
acaaggctcg acgcaaggcg ccacatttta ttcttcgaac agaattgattg ccattggcaa 840  
gaccttttga atttcgcaaa attagatttc aactcgttac aaaagttgca ccagagggag 900  
ctatgtgaaa tcacaagggtg gtggaaagat ctggattttg ccaagaaact accttttgcc 960  
agagacagaa tggtagagtg ctacttctgg atacttggg tgtactttga gcccataat 1020

## 497047kxr.ST25.txt

```

ttgcgtgcta ggaggatgct aaccaaggtg attgccttga cttccattat cgatgacatc 1080
tacgatgtct acggtacctt ggaagaactt gttctcttca ctgatgcaat tgagaggtgg 1140
gaaattagtg ccttggataa ccttcagat tataatgaaac tatgttatca agcacttttg 1200
gatgtttata gtatgattga tgaagagatg gccaaagcaag gaagatctta ttgcgtagac 1260
tatgcaaaat cttcaatgaa aatttttggtt agagcatact tcgaagaagc caaatgggtt 1320
caccaaggat atgttccaac tatggaagag tataatgcaag ttgcattagt aaccgcgggt 1380
tacaaaatgc ttgcaacctc ttcctttggtt ggcatgggag agttggcaac caaagaggcc 1440
tttgattggg tgtcaaatga tcctttaatt gttcaagctg catcagtgat aggcagactc 1500
aaggatgaca ttgttggcca caagtttgag caaaagagag ggcacgtggc gtcggctgtc 1560
gaatgctaca gtaagcaaca tggtacaata gaggaagagg ctattattga attggataaa 1620
caagttacac attcatggaa agacatcaac gcagagtgcc tctgcccaat caaggtccca 1680
atgcctcttc ttgcgcgagt tctcaatctt gcacgagtgc tttatgttat ataccaggat 1740
gaagacggct acactcattc tggaaccaag gtcaagaact ttgcaacctc agtgcttattc 1800
gattctatgc caatcaatta gaaaatgtaa caagacactg aagtggaggc ataaataaat 1860
tcaaaagttg gcttaaagtt gggctaaaaa aaaaaaa 1897

```

&lt;210&gt; 5

&lt;211&gt; 491

&lt;212&gt; PRT

&lt;213&gt; Actinidia deliciosa

&lt;400&gt; 5

```

Met Gln Leu Pro Cys Ala Gln Ala Leu Pro Ile Pro Thr Val Thr Thr
1          5          10          15

```

```

Asn Thr Ser Ile Glu Pro Pro His Val Thr Arg Arg Ser Ala Asn Tyr
          20          25          30

```

```

His Pro Ser Ile Trp Gly Asp His Phe Leu Ala Tyr Ser Ser Asp Ala
          35          40          45

```

```

Met Glu Glu Glu Asp Ile Asn Met Glu Gln Gln Gln Arg Leu His His
50          55          60

```

```

Leu Lys Gln Lys Val Arg Lys Met Leu Glu Ala Ala Ala Glu Gln Ser
65          70          75          80

```

497047kxr.ST25.txt

Ser Gln Met Leu Asn Leu Val Asp Lys Ile Gln Arg Leu Gly Val Ser  
                   85                  90                  95

Tyr His Phe Glu Thr Glu Ile Glu Thr Ala Leu Arg His Ile Tyr Lys  
                   100                  105                  110

Thr Cys Asp Tyr His Phe Asp Asp Leu His Thr Ala Ala Leu Ser Phe  
                   115                  120                  125

Arg Leu Leu Arg Gln Gln Gly Tyr Pro Val Ser Cys Asp Met Phe Asp  
                   130                  135                  140

Lys Phe Lys Asn Ser Lys Gly Glu Phe Gln Glu Ser Ile Ile Ser Asp  
                   145                  150                  155                  160

Val Arg Gly Met Leu Ser Leu Tyr Glu Ala Thr Cys Leu Met Ile His  
                   165                  170                  175

Gly Glu Asp Ile Leu Asp Glu Ala Leu Ala Phe Thr Ile Thr Gln Leu  
                   180                  185                  190

Arg Ser Ala Leu Pro Asn Leu Ser Thr Pro Phe Lys Glu Gln Ile Ile  
                   195                  200                  205

His Ala Leu Asn Gln Pro Ile His Lys Gly Leu Thr Arg Leu Asn Ala  
                   210                  215                  220

Arg Ser His Ile Leu Phe Phe Glu Gln Asn Asp Cys His Ser Lys Asp  
                   225                  230                  235                  240

Leu Leu Asn Phe Ala Lys Leu Asp Phe Asn Leu Leu Gln Lys Leu His  
                   245                  250                  255

Gln Arg Glu Leu Cys Glu Ile Thr Arg Glu Ile Ser Ala Leu Asp Gln  
                   260                  265                  270

Leu Pro Glu Tyr Met Lys Leu Cys Tyr Gln Ala Leu Leu Asp Val Tyr  
                   275                  280                  285

Ser Met Ile Asp Glu Glu Met Ala Lys Gln Gly Arg Ser Tyr Cys Val  
                   290                  295                  300

Asp Tyr Ala Lys Ser Ser Met Lys Ile Leu Val Arg Ala Tyr Phe Glu  
                   305                  310                  315                  320

Glu Ala Lys Trp Phe His Gln Gly Tyr Val Pro Thr Met Glu Glu Tyr  
                   325                  330                  335

497047kxr.ST25.txt

Met Gln Val Ala Leu Val Thr Ala Gly Tyr Lys Met Leu Ala Thr Ser  
 340 345 350

Ser Phe Val Gly Met Gly Glu Leu Ala Thr Lys Glu Ala Phe Asp Trp  
 355 360 365

Val Ser Asn Asp Pro Leu Ile Val Gln Ala Ala Ser Val Ile Gly Arg  
 370 375 380

Leu Lys Asp Asp Ile Val Gly His Lys Phe Glu Gln Lys Arg Gly His  
 385 390 395 400

Val Ala Ser Ala Val Glu Cys Tyr Ser Lys Gln His Gly Thr Thr Glu  
 405 410 415

Glu Glu Ala Ile Ile Glu Leu Tyr Lys Gln Val Thr His Ser Trp Lys  
 420 425 430

Asp Met Asn Ala Glu Cys Leu Cys Pro Thr Lys Val Pro Met Pro Leu  
 435 440 445

Leu Ala Arg Val Leu Asn Leu Ala Arg Val Leu Tyr Val Ile Tyr Gln  
 450 455 460

Asp Ala Asp Gly Tyr Thr His Ser Gly Thr Lys Val Lys Asn Phe Val  
 465 470 475 480

Thr Ser Val Leu Ile Asp Ser Met Pro Ile Asn  
 485 490

&lt;210&gt; 6

&lt;211&gt; 565

&lt;212&gt; PRT

&lt;213&gt; Actinidia deliciosa

&lt;400&gt; 6

Met Gln Leu Pro Cys Ala Gln Ala Leu Pro Ile Pro Thr Val Thr Thr  
 1 5 10 15

Asn Thr Ser Ile Glu Pro Pro His Val Thr Arg Arg Ser Ala Asn Tyr  
 20 25 30

His Pro Ser Ile Trp Gly Asp His Phe Leu Ala Tyr Ser Ser Asp Ala  
 35 40 45

497047kxr.ST25.txt

Met Glu Glu Glu Asp Ile Asn Met Glu Gln Gln Gln Arg Leu His His  
 50 55 60

Leu Lys Gln Lys Val Arg Lys Met Leu Glu Ala Ala Ala Lys Gln Ser  
 65 70 75 80

Ser Gln Met Leu Asn Leu Val Asp Lys Ile Gln Arg Leu Gly Val Ser  
 85 90 95

Tyr His Phe Glu Thr Glu Ile Glu Thr Ala Leu Arg His Ile Tyr Lys  
 100 105 110

Thr Cys Asp Tyr His Phe Asp Asp Leu His Thr Ala Ala Leu Ser Phe  
 115 120 125

Arg Leu Leu Arg Gln Gln Gly Tyr Pro Val Ser Cys Asp Met Phe Gly  
 130 135 140

Lys Phe Lys Asn Cys Lys Gly Glu Phe Gln Glu Ser Ile Ile Ser Asp  
 145 150 155 160

Val Arg Gly Met Leu Ser Leu Tyr Glu Ala Thr Cys Leu Arg Ile Arg  
 165 170 175

Gly Glu Asp Ile Leu Asp Glu Ala Leu Ala Phe Thr Thr Thr Gln Leu  
 180 185 190

Gln Ser Ala Leu Pro Asn Leu Ser Thr Pro Ile Lys Glu Gln Ile Ile  
 195 200 205

His Ala Leu Asn Gln Pro Ile His Lys Trp Leu Thr Arg Leu Asp Ala  
 210 215 220

Arg Arg His Ile Leu Phe Phe Glu Gln Asn Asp Cys His Gly Lys Asp  
 225 230 235 240

Leu Leu Asn Phe Ala Lys Leu Asp Phe Asn Ser Leu Gln Lys Leu His  
 245 250 255

Gln Arg Glu Leu Cys Glu Ile Thr Arg Trp Trp Lys Asp Leu Asp Phe  
 260 265 270

Ala Lys Lys Leu Pro Phe Ala Arg Asp Arg Met Val Glu Cys Tyr Phe  
 275 280 285

Trp Ile Leu Gly Val Tyr Phe Glu Pro Gln Tyr Leu Arg Ala Arg Arg  
 290 295 300

## 497047kxr.ST25.txt

Met Leu Thr Lys Val Ile Ala Leu Thr Ser Ile Ile Asp Asp Ile Tyr  
 305 310 315 320

Asp Val Tyr Gly Thr Leu Glu Glu Leu Val Leu Phe Thr Asp Ala Ile  
 325 330 335

Glu Arg Trp Glu Ile Ser Ala Leu Asp Asn Leu Pro Asp Tyr Met Lys  
 340 345 350

Leu Cys Tyr Gln Ala Leu Leu Asp Val Tyr Ser Met Ile Asp Glu Glu  
 355 360 365

Met Ala Lys Gln Gly Arg Ser Tyr Cys Val Asp Tyr Ala Lys Ser Ser  
 370 375 380

Met Lys Ile Leu Val Arg Ala Tyr Phe Glu Glu Ala Lys Trp Phe His  
 385 390 395 400

Gln Gly Tyr Val Pro Thr Met Glu Glu Tyr Met Gln Val Ala Leu Val  
 405 410 415

Thr Ala Gly Tyr Lys Met Leu Ala Thr Ser Ser Phe Val Gly Met Gly  
 420 425 430

Glu Leu Ala Thr Lys Glu Ala Phe Asp Trp Val Ser Asn Asp Pro Leu  
 435 440 445

Ile Val Gln Ala Ala Ser Val Ile Gly Arg Leu Lys Asp Asp Ile Val  
 450 455 460

Gly His Lys Phe Glu Gln Lys Arg Gly His Val Ala Ser Ala Val Glu  
 465 470 475 480

Cys Tyr Ser Lys Gln His Gly Thr Ile Glu Glu Glu Ala Ile Ile Glu  
 485 490 495

Leu Asp Lys Gln Val Thr His Ser Trp Lys Asp Ile Asn Ala Glu Cys  
 500 505 510

Leu Cys Pro Ile Lys Val Pro Met Pro Leu Leu Ala Arg Val Leu Asn  
 515 520 525

Leu Ala Arg Val Leu Tyr Val Ile Tyr Gln Asp Glu Asp Gly Tyr Thr  
 530 535 540

His Ser Gly Thr Lys Val Lys Asn Phe Ala Thr Ser Val Leu Ile Asp  
 545 550 555 560

497047kxr.ST25.txt

Ser Met Pro Ile Asn  
565

&lt;210&gt; 7

&lt;211&gt; 686

&lt;212&gt; DNA

&lt;213&gt; Actinidia chinensis

&lt;400&gt; 7

```

atcttattgc gtagactatg caaaatcttc aatgaaaagt ttggttagag catacttoga      60
agaagccaaa tggtttcacc aaggatatgt tccaactatg gaagagtata tgcaagttgc      120
aatagtaacc ggggcttaca aaattcttgc aaccacttcc tttgttggca tgggagagtt      180
ggcaacccaaa gaggtctttg attgggtgtc aaatgatcct ttaattgttc aagctgcac      240
aattgtttcc agactcacgg atgacattgt tggccacaag tttgagcaaa agagagggca      300
cgtggcatcg gcggttgaat gctacatgaa gcaacatggt acaacagagg aagaggccat      360
tgttgaattg tataagcaag ttacaaatgc atggaaagac atgaatgcag agtgcctctt      420
ccccaccaag gtcccaatgc ctcttctcgt gagagtcttc aatcttgcac gagtgattaa      480
tgttctatac aaggatgaag atggctacac tcattcaaga accaagggtta agaaatttgt      540
gacctcagtg cttgtagatt ttgtgccgat cagctagcaa acgttctctt ctaccacatg      600
ttaattagtc tgcttgctaa tgcagtttac taatatgaaa tttataaat gcgtattttc      660
caataaagga atttaaaaaa aaaaaa      686

```

&lt;210&gt; 8

&lt;211&gt; 190

&lt;212&gt; PRT

&lt;213&gt; Actinidia chinensis

&lt;400&gt; 8

```

Tyr Cys Val Asp Tyr Ala Lys Ser Ser Met Lys Ser Leu Val Arg Ala
1          5          10          15

Tyr Phe Glu Glu Ala Lys Trp Phe His Gln Gly Tyr Val Pro Thr Met
20          25          30

Glu Glu Tyr Met Gln Val Ala Ile Val Thr Gly Ala Tyr Lys Ile Leu
35          40          45

```



497047kxr.ST25.txt

Ala Thr Thr Ser Phe Val Gly Met Gly Glu Leu Ala Thr Lys Glu Val  
 50 55 60

Phe Asp Trp Val Ser Asn Asp Pro Leu Ile Val Gln Ala Ala Ser Ile  
 65 70 75 80

Val Ser Arg Leu Thr Asp Asp Ile Val Gly His Lys Phe Glu Gln Lys  
 85 90 95

Arg Gly His Val Ala Ser Ala Val Glu Cys Tyr Met Lys Gln His Gly  
 100 105 110

Thr Thr Glu Glu Glu Ala Ile Val Glu Leu Tyr Lys Gln Val Thr Asn  
 115 120 125

Ala Trp Lys Asp Met Asn Ala Glu Cys Leu Phe Pro Thr Lys Val Pro  
 130 135 140

Met Pro Leu Leu Val Arg Val Leu Asn Leu Ala Arg Val Ile Asn Val  
 145 150 155 160

Leu Tyr Lys Asp Glu Asp Gly Tyr Thr His Ser Arg Thr Lys Val Lys  
 165 170 175

Lys Phe Val Thr Ser Val Leu Val Asp Phe Val Pro Ile Ser  
 180 185 190

&lt;210&gt; 9

&lt;211&gt; 755

&lt;212&gt; DNA

&lt;213&gt; Vaccinium corymbosum

&lt;400&gt; 9

ggaagccaaa tggtttcatg aaggttatgt tccgagtatg gaagagtata tgagagttgc 60  
 actgggttacc ggtgcttaca aaatgcttgc aaccacttct tttgttggca tgggggattt 120  
 ggtgacccaaa gaggcctttg aatgggtgtc aagtgatect ttaattgttg aagctgcac 180  
 cgtgatttgc agactcatgg atgatatggc aggccacaag tttgagcaag agagaggaca 240  
 cgtggcttcg gcagttgaat gctacatgaa acaacatggt gcaacacaag aagtggttct 300  
 tcttgaattt aaaaaaagag ttacaaatgc atggaaagac atgaacgcag agtgcctccg 360  
 cccaactgcc gttccaatgc ctctctcac ccgagttctc aatctcgac gagtgatcaa 420  
 tgttatatac aaggatgaag atgggtacac tcattctgga acaaagctca agaactttgt 480  
 aatctcagtg cttatcgatt ctgtgccgat caattagcaa acagtagtcc taacttaa 540

## 497047kxr.ST25.txt

```

aatctgttgg cttataactt tataagtgtc gtgaaatgtt ctagtgaact tggtaaggat      600
gtattttccga tatgtagctc tatctccact gtacggttgt aatcttgctc tcttctacta      660
agaaagctca ttaatcgctg cttaaaatgt aaagccaact tgctcaagtt tatcgtcaaa      720
caagttctgt tttagcattt ttgttgga aa      755

```

&lt;210&gt; 10

&lt;211&gt; 171

&lt;212&gt; PRT

<213> *Vaccinium corymbosum*

&lt;400&gt; 10

```

Glu Ala Lys Trp Phe His Glu Gly Tyr Val Pro Ser Met Glu Glu Tyr
1          5          10          15

```

```

Met Arg Val Ala Leu Val Thr Gly Ala Tyr Lys Met Leu Ala Thr Thr
          20          25          30

```

```

Ser Phe Val Gly Met Gly Asp Leu Val Thr Lys Glu Ala Phe Glu Trp
          35          40          45

```

```

Val Ser Ser Asp Pro Leu Ile Val Glu Ala Ala Ser Val Ile Cys Arg
          50          55          60

```

```

Leu Met Asp Asp Met Ala Gly His Lys Phe Glu Gln Glu Arg Gly His
65          70          75          80

```

```

Val Ala Ser Ala Val Glu Cys Tyr Met Lys Gln His Gly Ala Thr Gln
          85          90          95

```

```

Glu Val Val Leu Leu Glu Phe Lys Lys Arg Val Thr Asn Ala Trp Lys
          100          105          110

```

```

Asp Met Asn Ala Glu Cys Leu Arg Pro Thr Ala Val Pro Met Pro Leu
          115          120          125

```

```

Leu Thr Arg Val Leu Asn Leu Ala Arg Val Ile Asn Val Ile Tyr Lys
          130          135          140

```

```

Asp Glu Asp Gly Tyr Thr His Ser Gly Thr Lys Leu Lys Asn Phe Val
145          150          155          160

```

```

Ile Ser Val Leu Ile Asp Ser Val Pro Ile Asn
          165          170

```

497047kxr.ST25.txt

&lt;210&gt; 11

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Synthetic primer

&lt;400&gt; 11

gaattccaac taccttgtgc tcaagc

26

&lt;210&gt; 12

&lt;211&gt; 25

&lt;212&gt; DNA

&lt;213&gt; Synthetic primer

&lt;400&gt; 12

ctcgagcctc cacttcagtg tcttg

25

&lt;210&gt; 13

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Synthetic primer

&lt;400&gt; 13

caattgagag gtgggagatc

20

&lt;210&gt; 14

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Synthetic primer

&lt;400&gt; 14

gttggaacat atccttggtg

20

&lt;210&gt; 15

&lt;211&gt; 20

&lt;212&gt; DNA

497047kxr.ST25.txt

&lt;213&gt; Synthetic primer

<400> 15  
taggcgtgtc ttaccat

20

&lt;210&gt; 16

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Synthetic primer

<400> 16  
gcatgaatga tttgttcctt

20

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ2003/000294

**A. CLASSIFICATION OF SUBJECT MATTER**Int. Cl. <sup>7</sup>: C07K 14/415; C07H 21/04; C12N 15/63, 15/82.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

See electronic database consulted below.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN File DGENE: Blastn (Seq. Id. No. 1) and Blastp (Seq. Id. No. 2). ANGIS: Blastn (Seq. Id. No.s 1, 3, 4 and 7) and Blastp (Seq. Id. No. 2).

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

| Category* | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                                                                                                            | Relevant to claim No. |
|-----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| P, X      | STEINMETZ, A. A. et al., NCBI Entrez Nucleotide, Accession Number AY561842 (online) 21 March 2004 (retrieved on 6 April 2004). Retrieved from the Internet: <URL: <a href="http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?db=nucleotide&amp;val=45478052">http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?db=nucleotide&amp;val=45478052</a> > (see sequences) | 1-44                  |
| P, X      | STEINMETZ, A. A. et al., NCBI Entrez Nucleotide, Accession Number AY561843 (online) 21 March 2004 (retrieved on 6 April 2004). Retrieved from the Internet: <URL: <a href="http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?db=nucleotide&amp;val=45478054">http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?db=nucleotide&amp;val=45478054</a> > (see sequences) | 1-44                  |

☒ Further documents are listed in the continuation of Box C☐ See patent family annex

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search  
6 April 2004

Date of mailing of the international search report

- 8 APR 2004

Name and mailing address of the ISA/AU

AUSTRALIAN PATENT OFFICE  
PO BOX 200, WODEN ACT 2606, AUSTRALIA  
E-mail address: pct@ipaustalia.gov.au  
Facsimile No. (02) 6285 3929

Authorized officer

MARIE-ANNE FAM

Telephone No : (02) 6283 2254

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ2003/000294

| C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT |                                                                                                                                                                                                                                                                                                                                                                      |                       |
|-------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| Category*                                             | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                                                                                                                   | Relevant to claim No. |
| X                                                     | VAN DER HOEVEN, R. S. et al., Genetic Control and Evolution of Sesquiterpene Biosynthesis in <i>Lycopersicon esculentum</i> and <i>L. hirsutum</i> .<br>The Plant Cell (2000), volume 12, pages 2283-2294<br>(see in particular page 2293, GenBank accession number AF279456, nucleotides 752-772)                                                                   | 37.                   |
| X                                                     | MARUYAMA, T. et al., NCBI Entrez Nucleotide, Accession Number AF282875 (online) 25 July 2001 (retrieved on 6 April 2004). Retrieved from the Internet: <URL: <a href="http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?db=nucleotide&amp;val=14717382">http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?db=nucleotide&amp;val=14717382</a> > (see nucleotides 1183-1199) | 37                    |

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ2003/000294

**Box I** Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos :  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos : 1, 13, 33 and 37 (in part)  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
Claims 1, 13 and 33 relate to any multifunctional germacrene-D synthase, however the description only provides support for particular proteins. Consequently the search for these claims was based on the germacrene-D synthase enzyme as exemplified in the specification.  
(continued on supplemental sheet)
3. ☐ Claims Nos :  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

**Box II** Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ2003/000294

**Supplemental Box**

(To be used when the space in any of Boxes I to VIII is not sufficient)

**Continuation of Box No: I**

Claim 37 defines any polynucleotide having at least 15 contiguous nucleotides from Seq. Id. No. 1. Thus the claim encompasses a vast number of possible sequences, particularly in view of the fact that the polynucleotide need not have any activity or function associated with germacrene-D synthase. A complete search of the claim was therefore not economically feasible. It is also noted that the documents cited against this claim represent only a small portion of the relevant prior art.